



WINTER **SPRING** **SUMMER** **AUTUMN**

HERBICIDE - Apply aminopyralid (Milestone™) before seedling emergence to basal rosette stage. **DO NOT APPLY MILESTONE UNTIL AFTER GRASS SEED HAS GERMINATED.** Apply at low rates - 3-5 fl. oz./acre.

SEEDING - Drill or broadcast seed annual ryegrass to fertilized pastures at 40-60 lbs N /acre.

HERBICIDE - Apply clopyralid (Transline™) from seedling emergence to basal rosette to bolting stage. Apply at low rates - 10-20 fl. oz./acre.

After plants visible, susceptible to hand pulling. Leave in field to decompose before flowering stage, after flowering stage, bag and dispose of plant in appropriate location.

MOWING - Mow plants with hand-held string trimmer after full bolt, early flowering.

HERBICIDE - apply glyphosate herbicides (Roundup™) at 1-2 quarts/acre at full bolt.

***CAUTION* SEED PRODUCTION STAGE**
PULL AND BAG FOR DISPOSAL - MOWING OR PULLING AT THIS STAGE WILL ALLOW SEED TO RIPEN IF LEFT ON THE GROUND

ORGANIC - FERTILIZATION + OVERSEEDING
 Apply organic fertilizers high in N and P in late fall before rains to prepare for annual grass seeding in the winter.

ORGANIC

HERBICIDE

LIFE STAGE

NOTES