



**WEEKLY SOIL MOISTURE LOSS IN INCHES**

(Estimated Crop Evapotranspiration or ET<sub>c</sub>)

04/20/18 through 04/26/18

Crops (Leafout Date)	#188 Madera II			#39 Parlier			#86 Lindcove		
	4/20 - 4/26 Water Use	Accum'd Seasonal Water Use	4/27 - 5/3 Estimated ETc	4/20 - 4/26 Water Use	Accum'd Seasonal Water Use	4/27 - 5/3 Estimated ETc	4/20 - 4/26 Water Use	Accum'd Seasonal Water Use	4/27 - 5/3 Estimated ETc
Almonds (3/16) *	1.09	4.08	1.10	1.11	4.19	1.07	1.12	3.99	1.00
Pistachio (4/21) * **	0.09	0.09	0.13	0.10	0.10	0.13	0.10	0.10	0.13
Citrus (2/1)	1.02	6.58	1.03	1.04	6.68	1.00	1.05	6.51	0.93
Raisin Grapes (3/16 ) (11 ft. row spacing) ***	0.36	0.84	0.42	0.37	0.88	0.42	0.37	0.82	0.42
Winegrapes (3/16 ) (10 ft. spacing on California Sprawl Trellis) ***	0.43	1.23	0.49	0.44	1.23	0.49	0.45	1.18	0.44
Walnuts (4/4)	0.79	1.27	0.83	0.80	1.29	0.83	0.80	1.24	0.78
Stone Fruit (3/16 )	0.58	1.86	0.65	0.59	1.89	0.65	0.59	1.82	0.60
Past 7 days precipitation (inches)	0.00			0.00			0.00		
Accumulated precipitation (inches) (1/1/2018)	6.33			4.96			3.32		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

\* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

\*\* Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

\*\*\* Raisin Grapes and Winegrapes Irrigation should hold off until midday leaf water potential drops to -1.0 MPa, before that soil moisture reservoir is sufficient to supply the vine water demand. Update will be sent shortly once the county wide leaf water potential reaches approximate -1.0 MPa. Growers should adjust the irrigation start date based on the individual vineyard location and soil type.

**PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY <sup>1</sup>**

Crops	#188 Madera II				#39 Parlier				#86 Lindcove			
	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/16)	1.7	1.5	1.3	1.1	1.7	1.5	1.3	1.2	1.7	1.5	1.3	1.2
Pistachio (4/21)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Citrus (2/1)	1.6	1.4	1.2	1.1	1.6	1.4	1.2	1.1	1.6	1.4	1.2	1.1
Raisin Grapes (3/16 ) (late season table, 75% cover)***	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Winegrapes (3/16 ) (10 ft. spacing on California Sprawl Trellis) ***	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Walnuts (4/4)	1.2	1.1	0.9	0.8	1.2	1.1	0.9	0.8	1.2	1.1	0.9	0.8
Stone Fruit (3/16 )	0.9	0.8	0.7	0.6	0.9	0.8	0.7	0.6	0.9	0.8	0.7	0.6

<sup>1</sup> The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

**PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE**

Crops	#188 Madera II				#39 Parlier				#86 Lindcove			
	401	354	307	260	401	354	307	283	401	354	307	283
Almonds 115 Trees/A	401	354	307	260	401	354	307	283	401	354	307	283
Pistachio 106 Trees/A	25	25	25	25	25	25	25	25	25	25	25	25
Citrus 110 Trees/A	395	346	296	272	395	346	296	272	395	346	296	272
Raisin Grapes 566 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Winegrapes 622 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Walnuts 76 Trees/A	429	393	322	286	429	393	322	286	429	393	322	286
Stonefruit 172 Trees/A	142	126	111	95	142	126	111	95	142	126	111	95

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.

**WEEKLY SOIL MOISTURE LOSS IN INCHES**  
 (Estimated Crop Evapotranspiration or ET<sub>c</sub>)  
 04/20/18 through 04/26/18

Crops (Leafout Date)	#124 Panoche			#2 Five Points			#15 Stratford		
	4/20 - 4/26 Water Use	Accum'd Seasonal Water Use	4/27 - 5/3 Estimated ETc	4/20 - 4/26 Water Use	Accum'd Seasonal Water Use	4/27 - 5/3 Estimated ETc	4/20 - 4/26 Water Use	Accum'd Seasonal Water Use	4/27 - 5/3 Estimated ETc
Almonds (3/16) *	1.18	4.45	1.22	1.30	5.03	1.28	1.32	5.05	1.24
Pistachio (4/21) * **	0.12	0.12	0.14	0.12	0.12	0.16	0.12	0.12	0.14
Citrus (2/1)	1.11	7.98	1.15	1.23	8.97	1.21	1.25	8.84	1.17
Raisin Grapes (3/16 ) (11 ft. row spacing) ***	0.36	0.87	0.45	0.41	0.98	0.49	0.41	0.98	0.47
Winegrapes (3/16 ) (10 ft. spacing on California Sprawl Trellis) ***	0.45	1.30	0.52	0.49	1.47	0.56	0.50	1.47	0.54
Walnuts (4/4)	0.85	1.35	0.93	0.91	1.52	0.99	0.93	1.55	0.95
Stone Fruit (3/16 )	0.63	2.05	0.75	0.69	2.31	0.75	0.70	2.30	0.75
<b>Past 7 days precipitation (inches)</b>	0.00			0.00			0.00		
<b>Accumulated precipitation (inches) (1/1/18)</b>	2.56			3.00			1.62		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

\* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

\*\* Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

\*\*\* Raisin Grapes and Winegrapes Irrigation should hold off until midday leaf water potential drops to -1.0 MPa, before that soil moisture reservoir is sufficient to supply the vine water demand. Update will be sent shortly once the county wide leaf water potential reaches approximate -1.0 MPa. Growers should adjust the irrigation start date based on the individual vineyard location and soil type.

**PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY <sup>1</sup>**

Crops	#124 Panoche				#2 Five Points				#15 Stratford			
	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/16)	1.8	1.6	1.4	1.2	2.0	1.7	1.5	1.4	2.0	1.8	1.6	1.4
Pistachio (4/21)	0.2	0.2	0.1	0.1	0.2	0.2	0.1	0.1	0.2	0.2	0.1	0.1
Citrus (2/1)	1.7	1.5	1.3	1.2	1.9	1.6	1.4	1.3	1.9	1.7	1.5	1.3
Raisin Grapes (3/16 ) (late season table, 75% cover)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Winegrapes (3/16 ) (10 ft. spacing on California Sprawl Trellis)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Walnuts (4/4)	1.3	1.1	1.0	0.9	1.4	1.2	1.1	1.0	1.4	1.2	1.1	1.0
Stone Fruit (3/16 )	1.0	0.8	0.7	0.7	1.1	0.9	0.8	0.7	1.1	0.9	0.8	0.7

<sup>1</sup> The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

**PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE**

Crops	#124 Panoche				#2 Five Points				#15 Stratford			
	425	378	331	283	472	401	354	331	472	425	378	331
Almonds 115 Trees/A	425	378	331	283	472	401	354	331	472	425	378	331
Pistachio 106 Trees/A	50	50	25	25	50	50	25	25	50	50	25	25
Citrus 110 Trees/A	420	370	321	296	469	395	346	321	469	420	370	321
Raisin Grapes 566 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Winegrapes 622 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Walnuts 76 Trees/A	464	393	357	322	500	429	393	357	500	429	393	357
Stonefruit 172 Trees/A	158	126	111	111	174	142	126	111	174	142	126	111

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.