HEALTHY FOOD SYSTEMS .

ALTHY ENVIRONMENT

University of California

Agriculture and Natural Resources



UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

WEEKLY SOIL MOISTURE LOSS IN INCHES (Estimated Crop Evapotranspiration or ET_c) 05/11/18 through 05/17/18

Crops (Leafout Date)	#1	#188 Madera II			#39 Parlier				#86 Lindcove		
	5/11 - 5/17	Accum'd	5/18 - 5/24		5/11 - 5/17	Accum'd	5/18 - 5/24		5/11 - 5/17	Accum'd	5/18 - 5/24
	Water	Seasonal	Estimated		Water	Seasonal	Estimated		Water	Seasonal	Estimated
	Use	Water Use	ETc		Use	Water Use	ETc		Use	Water Use	ETc
Almonds (3/16) *	1.43	7.94	1.49		1.38	7.94	1.42		1.35	7.52	1.36
Pistachio (4/21) * **	0.50	1.05	0.67		0.46	0.99	0.63		0.47	0.97	0.61
Citrus (2/1)	1.22	10.00	1.19		1.17	9.96	1.14		1.14	9.58	1.08
Raisin Grapes (3/16) (11 ft. row spacing)	1.04	3.23	1.14		1.01	3.21	1.07		0.97	2.99	1.05
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	0.95	3.50	1.05		0.93	3.44	1.00		0.90	3.27	0.94
Walnuts (4/4)	1.16	4.40	1.35		1.12	4.33	1.28		1.10	4.12	1.22
Stone Fruit (3/16)	0.97	4.37	1.14		0.94	4.31	1.07		0.92	4.14	1.05
Past 7 days precipitation (inches)		0.00				0.00				0.00	
Accumulated precipitation (inches) (1/1/2018)		6.33				4.96				3.32	

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

P	AST WEEK	LY APPLI	ED WATEI	R IN INCHE	ES, ADJUST	FED FOR E	FFICIENC	CY 1				
Crops		#188 Made		#39 Parlier								
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/16)	2.2	1.9	1.7	1.5	2.1	1.8	1.6	1.5	2.1	1.8	1.6	1.4
Pistachio (4/21)	0.8	0.7	0.6	0.5	0.8	0.7	0.6	0.5	0.8	0.7	0.6	0.5
Citrus (2/1)	1.9	1.6	1.4	1.3	1.8	1.6	1.4	1.2	1.8	1.5	1.3	1.2
Raisin Grapes (3/16) (11 ft. row spacing)	1.6	1.4	1.2	1.1	1.6	1.3	1.2	1.1	1.5	1.3	1.1	1.0
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	1.5	1.3	1.1	1.0	1.4	1.2	1.1	1.0	1.4	1.2	1.1	0.9
Walnuts (4/4)	1.8	1.5	1.4	1.2	1.7	1.5	1.3	1.2	1.7	1.5	1.3	1.2
Stone Fruit (3/16)	1.5	1.3	1.1	1.0	1.4	1.3	1.1	1.0	1.4	1.2	1.1	1.0

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST V	VEEKLY A	PPLIED W	ATER IN G	FALLON PI	ER TREE O	R VINE					
Crops	#188 Madera II					#39 Parlier						
Almonds 115 Trees/A	519	449	401	354	496	425	378	354	496	425	378	331
Pistachio 106 Trees/A	199	174	149	125	199	174	149	125	199	174	149	125
Citrus 110 Trees/A	469	395	346	321	444	395	346	296	444	370	321	296
Raisin Grapes 566 Vines/A	77	67	58	53	77	62	58	53	72	62	53	48
Winegrapes 622 Vines/A	65	57	48	44	61	52	48	44	61	52	48	39
Walnuts 76 Trees/A	643	536	500	429	607	536	464	429	607	536	464	429
Stonefruit 172 Trees/A	237	205	174	158	221	205	174	158	221	189	174	158
For further information concerning all counties receiving this report, contact	the Fresno (Co. Farm Ad	visor's office	at (559) 24	1-7526.							

University of California

HEALTHY FOOD SYSTEMS

Agriculture and Natural Resources



UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

		nated Crop	OISTURE Evapotrans 8 through 0	spiration or							
Crops (Leafout Date)	#	124 Panoch	ie		#.	2 Five Poin	ts	#	15 Stratfor	d	
	5/11 - 5/17	Accum'd	5/18 - 5/24		5/11 - 5/17	Accum'd	5/18 - 5/24	5/11 - 5/17	Accum'd	5/18 - 5/24	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (3/16) *	1.47	8.30	1.54		1.52	9.24	1.61	1.53	9.22	1.62	
Pistachio (4/21) * **	0.50	1.05	0.68		0.53	1.15	0.74	0.53	1.14	0.74	
Citrus (2/1)	1.25	11.36	1.26		1.28	12.66	1.32	1.29	12.48	1.33	
Raisin Grapes (3/16) (11 ft. row spacing)	1.04	3.14	1.12		1.06	3.42	1.19	1.06	3.40	1.20	
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	0.94	3.46	1.05		0.97	3.80	1.11	0.98	3.81	1.12	
Walnuts (4/4)	1.19	4.46	1.47		1.25	4.94	1.48	1.25	4.94	1.48	
Stone Fruit (3/16)	1.01	4.57	1.19		1.03	5.02	1.25	1.02	4.97	1.26	
Past 7 days precipitation (inches)		0.00				0.00			0.00		
Accumulated precipitation (inches) (1/1/2018)		2.56				3.04			1.62		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

Р	AST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹												
Crops	#124 Panoche					#2 Five Poi	ints						
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/16)	2.3	2.0	1.7	1.5	2.3	2.0	1.8	1.6	2.4	2.0	1.8	1.6	
Pistachio (4/21)	0.8	0.7	0.6	0.5	0.8	0.7	0.6	0.6	0.8	0.7	0.6	0.6	
Citrus (2/1)	1.9	1.7	1.5	1.3	2.0	1.7	1.5	1.3	2.0	1.7	1.5	1.4	
Raisin Grapes (3/16) (11 ft. row spacing)	1.6	1.4	1.2	1.1	1.6	1.4	1.2	1.1	1.6	1.4	1.2	1.1	
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	1.4	1.3	1.1	1.0	1.4	1.3	1.1	1.0	1.5	1.3	1.2	1.0	
Walnuts (4/4)	1.8	1.6	1.4	1.3	1.9	1.7	1.5	1.3	1.9	1.7	1.5	1.3	
Stone Fruit (3/16)	1.6	1.3	1.2	1.1	1.6	1.4	1.2	1.1	1.6	1.4	1.2	1.1	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST	WEEKLY A	PPLIED W	ATER IN (FALLON PL	ER TREE (DR VINE					
Crops		#124 Pano	che			#2 Five Poi	ints					
Almonds 115 Trees/A	543	472	401	354	543	472	425	378	567	472	425	378
Pistachio 106 Trees/A	199	174	149	125	199	174	149	149	199	174	149	149
Citrus 110 Trees/A	469	420	370	321	494	420	370	321	494	420	370	346
Raisin Grapes 566 Vines/A	77	67	58	53	77	67	58	53	77	67	58	53
Winegrapes 622 Vines/A	61	57	48	44	61	57	48	44	65	57	52	44
Walnuts 76 Trees/A	643	572	500	464	679	607	536	464	679	607	536	464
Stonefruit 172 Trees/A	253	205	189	174	253	221	189	174	253	221	189	174
For further information concerning all counties receiving this report.	contact the Fresno	Co Farm Ad	visor's office	e at (559) 24	1-7526							