University of California

Agriculture and Natural Resources



UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

WEEKLY SOIL MOISTURE LOSS IN INCHES (Estimated Crop Evapotranspiration or ET_C) 05/25/18 through 05/31/18

Crops (Leafout Date)	#1						#86 Lindcov	e		
	5/25 - 5/31	Accum'd	6/1 - 6/7		5/25 - 5/31	Accum'd	6/1 - 6/7	5/25 - 5/31	Accum'd	6/1 - 6/7
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc
Almonds (3/16) *	1.57	10.97	1.69		1.51	10.89	1.68	1.40	10.30	1.65
Pistachio (4/21) * **	0.86	2.56	1.09		0.83	2.47	1.03	0.79	2.37	1.00
Citrus (2/1)	1.17	12.34	1.20		1.12	12.25	1.19	1.05	11.74	1.16
Raisin Grapes (3/16) (11 ft. row spacing)	1.35	5.80	1.47		1.29	5.70	1.40	1.20	5.35	1.37
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	1.28	5.91	1.40		1.23	5.76	1.33	1.15	5.49	1.33
Walnuts (4/4)	1.37	7.09	1.52		1.31	6.96	1.50	1.22	6.59	1.47
Stone Fruit (3/16)	1.18	6.67	1.40		1.14	6.55	1.40	1.06	6.24	1.37
Past 7 days precipitation (inches)		0.00				0.00			0.00	
Accumulated precipitation (inches) (1/1/2018)		6.33				4.96			3.32	

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY 1												
Crops	#188 Madera II				#39 Parlier							
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/16)	2.4	2.1	1.8	1.7	2.3	2.0	1.8	1.6	2.2	1.9	1.6	1.5
Pistachio (4/21)	1.3	1.1	1.0	0.9	1.3	1.1	1.0	0.9	1.3	1.1	1.0	0.9
Citrus (2/1)	1.8	1.6	1.4	1.2	1.7	1.5	1.3	1.2	1.6	1.4	1.2	1.1
Raisin Grapes (3/16) (11 ft. row spacing)	2.1	1.8	1.6	1.4	2.0	1.7	1.5	1.4	1.8	1.6	1.4	1.3
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	2.0	1.7	1.5	1.3	1.9	1.6	1.4	1.3	1.8	1.5	1.4	1.2
Walnuts (4/4)	2.1	1.8	1.6	1.4	2.0	1.7	1.5	1.4	1.9	1.6	1.4	1.3
Stone Fruit (3/16)	1.8	1.6	1.4	1.2	1.8	1.5	1.3	1.2	1.6	1.4	1.2	1.1

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST V	VEEKLY A	PPLIED W	ATER IN G	ALLON P	ER TREE C	R VINE							
Crops		#188 Madera II #39 Parlier									#86 Lindcove			
Almonds 115 Trees/A	567	496	425	401	543	472	425	378	519	449	378	354		
Pistachio 106 Trees/A	324	274	249	224	324	274	249	224	324	274	249	224		
Citrus 110 Trees/A	444	395	346	296	420	370	321	296	395	346	296	272		
Raisin Grapes 566 Vines/A	101	86	77	67	96	82	72	67	86	77	67	62		
Winegrapes 622 Vines/A	87	74	65	57	83	70	61	57	79	65	61	52		
Walnuts 76 Trees/A	750	643	572	500	715	607	536	500	679	572	500	464		
Stonefruit 172 Trees/A	284	253	221	189	284	237	205	189	253	221	189	174		
For further information concerning all counties receiving this report, contac	t the Fresno (Co. Farm Ad	visor's office	at (559) 24	1-7526.									

UCCE/DWR Weekly Crop

Water Use Report

University of California

Agriculture and Natural Resources

Making a Difference for California					_						
	WEEKL	Y SOIL M	OISTURE	LOSS IN	INCHES						
	(Esti	mated Crop	-	-	ET _C)						
		05/25/1	8 through ()5/31/18							
Crops (Leafout Date)	#	#124 Panoch	e		#	2 Five Poin	ts	#	15 Stratfor	d	
	5/25 - 5/31	Accum'd	6/1 - 6/7		5/25 - 5/31	Accum'd	6/1 - 6/7	5/25 - 5/31	Accum'd	6/1 - 6/7	1
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated	1
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc	┝───
Almonds (3/16) *	1.63	11.46	1.82		1.64	12.43	1.88	1.62	12.42	1.89	ł
Pistachio (4/21) * **	0.90	2.63	1.15		0.91	2.75	1.16	0.89	2.74	1.16	ł
Citrus (2/1)	1.22	13.80	1.27		1.25	15.15	1.32	1.21	14.96	1.33	l
Raisin Grapes (3/16) (11 ft. row spacing)	1.34	5.67	1.53		1.36	6.01	1.59	1.35	6.00	1.60	ł
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	1.28	5.84	1.46		1.29	6.21	1.52	1.28	6.23	1.53	1
Walnuts (4/4)	1.41	7.27	1.63		1.44	7.82	1.69	1.42	7.82	1.70	ł
Stone Fruit (3/16)	1.21	6.95	1.53		1.22	7.43	1.59	1.21	7.40	1.60	
Past 7 days precipitation (inches)		0.00				0.00			0.00		
Accumulated precipitation (inches) (1/1/2018)		2.56				3.04			1.62		l

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

P	AST WEEK	LY APPLI	ED WATEH	R IN INCHI	ES, ADJUST	FED FOR E	FFICIENC	CY 1				
Crops	#124 Panoche					#2 Five Poi	ints					
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/16)	2.5	2.2	1.9	1.7	2.5	2.2	1.9	1.7	2.5	2.2	1.9	1.7
Pistachio (4/21)	1.4	1.2	1.1	0.9	1.4	1.2	1.1	1.0	1.4	1.2	1.0	0.9
Citrus (2/1)	1.9	1.6	1.4	1.3	1.9	1.7	1.5	1.3	1.9	1.6	1.4	1.3
Raisin Grapes (3/16) (11 ft. row spacing)	2.1	1.8	1.6	1.4	2.1	1.8	1.6	1.4	2.1	1.8	1.6	1.4
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	2.0	1.7	1.5	1.3	2.0	1.7	1.5	1.3	2.0	1.7	1.5	1.3
Walnuts (4/4)	2.2	1.9	1.7	1.5	2.2	1.9	1.7	1.5	2.2	1.9	1.7	1.5
Stone Fruit (3/16)	1.9	1.6	1.4	1.3	1.9	1.6	1.4	1.3	1.9	1.6	1.4	1.3

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%. DAGE WEEKLY, A DRIVED WATER IN GALLON DER TREE OR VINE

	PAST V	VEEKLY A	PPLIED W	ATER IN G	GALLON PL	ER TREE (OR VINE					
Crops		#124 Pano	che			#2 Five Poi	ints					
Almonds 115 Trees/A	590	519	449	401	590	519	449	401	590	519	449	401
Pistachio 106 Trees/A	349	299	274	224	349	299	274	249	349	299	249	224
Citrus 110 Trees/A	469	395	346	321	469	420	370	321	469	395	346	321
Raisin Grapes 566 Vines/A	101	86	77	67	101	86	77	67	101	86	77	67
Winegrapes 622 Vines/A	87	74	65	57	87	74	65	57	87	74	65	57
Walnuts 76 Trees/A	786	679	607	536	786	679	607	536	786	679	607	536
Stonefruit 172 Trees/A	300	253	221	205	300	253	221	205	300	253	221	205
For further information concerning all counties receiving this report, contact	the Fresno (Co. Farm Ad	visor's office	e at (559) 24	1-7526.				•			