LTHY ENVIRONMENT

S • HEALTHY COMMUNITIES • HEALTHY CALIFORNIAN

University of **California** Agriculture and Natural Resources



UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

WEEKLY SOIL MOISTURE LOSS IN INCHES (Estimated Crop Evapotranspiration or ET_C) 09/21/18 through 09/27/18

Crops (Leafout Date)	#18	#188 Madera II *** #39 Parlier #86 Lindcove					e			
	9/21-9/27	Accum'd	9/28-10/4		9/21-9/27	Accum'd	9/28-10/4	9/21-9/27	Accum'd	9/28-10/4
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc
Almonds (3/16) *	1.19	40.59	1.02		1.33	42.46	1.00	1.33	41.26	0.98
Pistachio (4/21) * **	1.26	31.36	1.02		1.40	33.32	1.00	1.40	32.70	0.99
Citrus (2/1)	0.75	31.32	0.66		0.85	32.48	0.64	0.84	31.48	0.63
Raisin Grapes (3/16) (11 ft. row spacing)	0.93	29.56	0.81		1.05	31.01	0.79	1.05	30.22	0.78
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	1.05	32.40	0.88		1.19	34.11	0.86	1.19	33.30	0.85
Walnuts (4/4)	1.10	36.24	0.91		1.21	38.20	0.89	1.21	37.29	0.89
Stone Fruit (3/16)	1.26	34.62	1.02		1.40	36.60	1.00	1.40	35.76	0.99
Past 7 days precipitation (inches)		0.00				0.00			0.00	
Accumulated precipitation (inches) (1/1/2018)		6.33				4.96			3.32	

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

*** CIMIS station #188 Madera II has been taken out of service due to a conversion of the pasture to permanent crops. For the remainder of 2018 irrigation season Historical Average ETo will be used for the weekly report.

P	AST WEEK	LY APPLI	ED WATEI	R IN INCHI	ES, ADJUST	FED FOR E	FFICIENC	Y ¹				
Crops	#188 Madera II				#39 Parlier							
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/16)	1.8	1.6	1.4	1.3	2.0	1.8	1.6	1.4	2.0	1.8	1.6	1.4
Pistachio (4/21)	1.9	1.7	1.5	1.3	2.2	1.9	1.6	1.5	2.2	1.9	1.6	1.5
Citrus (2/1)	1.2	1.0	0.9	0.8	1.3	1.1	1.0	0.9	1.3	1.1	1.0	0.9
Raisin Grapes (3/16) (11 ft. row spacing)	1.4	1.2	1.1	1.0	1.6	1.4	1.2	1.1	1.6	1.4	1.2	1.1
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	1.6	1.4	1.2	1.1	1.8	1.6	1.4	1.3	1.8	1.6	1.4	1.3
Walnuts (4/4)	1.7	1.5	1.3	1.2	1.9	1.6	1.4	1.3	1.9	1.6	1.4	1.3
Stone Fruit (3/16)	1.9	1.7	1.5	1.3	2.2	1.9	1.6	1.5	2.2	1.9	1.6	1.5

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST V	VEEKLY A	PPLIED W	ATER IN (GALLON PI	ER TREE C	R VINE						
Crops	#188 Madera II #39 Parlier									#86 Lindcove			
Almonds 115 Trees/A	425	378	331	307	472	425	378	331	472	425	378	331	
Pistachio 106 Trees/A	473	424	374	324	548	473	399	374	548	473	399	374	
Citrus 110 Trees/A	296	247	222	197	321	272	247	222	321	272	247	222	
Raisin Grapes 566 Vines/A	67	58	53	48	77	67	58	53	77	67	58	53	
Winegrapes 622 Vines/A	70	61	52	48	79	70	61	57	79	70	61	57	
Walnuts 76 Trees/A	607	536	464	429	679	572	500	464	679	572	500	464	
Stonefruit 172 Trees/A	300	268	237	205	347	300	253	237	347	300	253	237	
For further information concerning all counties receiving this report, contact	the Fresno (Co. Farm Ad	visor's office	at (559) 24	1-7526.				•				

HEALTHY FOOD SYSTEMS .

ALTHY ENVIRONMENTS

HEALTHY COMMUNITIES HEALTHY CALIFORNIAN

University of California

Agriculture and Natural Resources



UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

		mated Crop	OISTURE Evapotran 8 through 0	spiration or							
Crops (Leafout Date)	#	124 Panoch	ne		#	² Five Poin	ts	#15 Stratford			
	9/21-9/27	Accum'd	9/28-10/4		9/21-9/27	Accum'd	9/28-10/4	 9/21-9/27	Accum'd	9/28-10/4	1
	Water Use	Seasonal Water Use	Estimated ETc		Water Use	Seasonal Water Use	Estimated ETc	Water Use	Seasonal Water Use	Estimated ETc	1
Almonds (3/16) *	1.30	44.22	1.21		1.47	47.06	1.23	1.49	46.86	1.26	
Pistachio (4/21) * **	1.37	34.58	1.28		1.54	36.55	1.30	1.56	36.30	1.33	1
Citrus (2/1)	0.82	34.69	0.78		0.94	37.24	0.80	0.96	36.93	0.83	1
Raisin Grapes (3/16) (11 ft. row spacing)	1.02	31.78	0.93		1.18	33.64	0.95	1.18	33.48	0.98	i
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	1.16	35.08	1.07		1.33	37.17	1.09	1.35	37.02	1.12	Ì
Walnuts (4/4)	1.17	39.58	1.08		1.33	42.01	1.08	1.35	41.78	1.10	Ì
Stone Fruit (3/16)	1.37	38.09	1.28		1.54	40.42	1.30	1.56	40.15	1.33	
Past 7 days precipitation (inches)		0.00				0.00			0.00		
Accumulated precipitation (inches) (1/1/2018)		2.56				3.04			1.62		i -

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

PA	AST WEEK	LY APPLI	ED WATEF	R IN INCHE	ES. ADIUST	FED FOR F	FFICIENC	V ¹				
Crops	#124 Panoche					#2 Five Poi		-				
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/16)	2.0	1.7	1.5	1.4	2.3	2.0	1.7	1.5	2.3	2.0	1.8	1.6
Pistachio (4/21)	2.1	1.8	1.6	1.4	2.4	2.1	1.8	1.6	2.4	2.1	1.8	1.6
Citrus (2/1)	1.3	1.1	1.0	0.9	1.4	1.3	1.1	1.0	1.5	1.3	1.1	1.0
Raisin Grapes (3/16) (11 ft. row spacing)	1.6	1.4	1.2	1.1	1.8	1.6	1.4	1.2	1.8	1.6	1.4	1.2
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis)	1.8	1.5	1.4	1.2	2.0	1.8	1.6	1.4	2.1	1.8	1.6	1.4
Walnuts (4/4)	1.8	1.6	1.4	1.2	2.0	1.8	1.6	1.4	2.1	1.8	1.6	1.4
Stone Fruit (3/16)	2.1	1.8	1.6	1.4	2.4	2.1	1.8	1.6	2.4	2.1	1.8	1.6

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST V	VEEKLY A	PPLIED W	ATER IN G	GALLON PI	ER TREE (OR VINE					
Crops		#124 Pano	che			#2 Five Poi	ints					
Almonds 115 Trees/A	472	401	354	331	543	472	401	354	543	472	425	378
Pistachio 106 Trees/A	523	448	399	349	598	523	448	399	598	523	448	399
Citrus 110 Trees/A	321	272	247	222	346	321	272	247	370	321	272	247
Raisin Grapes 566 Vines/A	77	67	58	53	86	77	67	58	86	77	67	58
Winegrapes 622 Vines/A	79	65	61	52	87	79	70	61	92	79	70	61
Walnuts 76 Trees/A	643	572	500	429	715	643	572	500	750	643	572	500
Stonefruit 172 Trees/A	332	284	253	221	379	332	284	253	379	332	284	253
For further information concerning all counties receiving this report, contac	t the Fresno	Co. Farm Ad	visor's office	at (559) 24	1-7526.				-			