

Agronomy of small grain production in California's North Coast region



Mark Lundy

UCCE Agronomy Advisor


Colusa-Sutter-Yuba Counties

- General agronomic concepts
- Pre-season considerations
- In-season considerations
- Resources

Liebig's law of the minimum

Justus von Liebig's
"Law of the Minimum"
published in 1873

"If one growth factor/nutrient is deficient, plant growth is limited, even if all other vital factors/nutrients are adequate...plant growth is improved by increasing the supply of the deficient factor/nutrient"



JUSTUS VON LIEBIG 1803 - 1873

<http://kemnovation.com/crop-nutrition/>





Figure 1-10 Liebig's Law of the Minimum states that the most limiting factor determines yield potential. Producers should minimize or eliminate the most limiting factor first, then the second most limiting factor, and so forth. Only in this manner can maximum yield potential be achieved (Source: Potash and Phosphate Institute).

Site selection



4/20/2013

Site selection



Crop history?

Apparent within-site
variability?

Precipitation?

Evapotranspiration?


Irrigation?

Slope?

Soil type?

3/13/2010

Rainfall and plant water demand



The screenshot shows the CIMIS website interface. At the top, there is a navigation bar with tabs for HOME, STATIONS, DATA, SPATIAL, and RESOURCES. Below this, there are sub-tabs for Station List, Station Location Map, Siting, Sensors, and Maintenance. A notice on the left states that the CIMIS ET-XML service will be discontinued. The main content area features a Bing Map showing station coordinate points. The map includes labels for roads like Star Rd, Brook Rd S, and Shiloh Rd, and landmarks like Charles M. Schulz Sonoma Co. Airport. The map is set to 'Bird's eye' view.

www.cimis.water.ca.gov/Stations.aspx

Login | Register

CIMIS

CALIFORNIA IRRIGATION MANAGEMENT INFORMATION SYSTEM
CALIFORNIA DEPARTMENT OF WATER RESOURCES

HOME STATIONS DATA SPATIAL RESOURCES

Notices

The CIMIS ET-XML service will soon be discontinued. FTP service will be changing in the near future.

See the System News for more details.

Station List Station Location Map Siting Sensors Maintenance

This Bing Map shows CIMIS station coordinate points. You can zoom in and out to see the exact station locations. Click the station marker for more detailed information.


Irrigate like a Pro

<http://www.cimis.water.ca.gov/>



Rainfall


Windsor, CA



Average rainfall total: 30.4 inches

Rainfall and evapotranspiration


Windsor, CA



$$\text{Aridity Index} [\text{Precipitation (30.4)} / \text{Evapotranspiration (37.9)}] = 0.80$$

Rainfall distribution & totals vary from year to year


Windsor, CA



2013-2014 rainfall total: 19.3 inches

Rainfall distribution & totals vary from year to year

Windsor, CA




2013-2014 rainfall total: 19.3 inches

2012-2013 rainfall total: 27.5 inches

- Small grains use 17 – 26 inches of soil water
 - amount that leaves the system via evapotranspiration
 - barley < oats < wheat
- Amount of water available to the crop will depend on:
 - what you grow
 - when you plant
 - irrigation?
 - SOIL
- How much will soil water contribute to total evapotranspiration?

How much of evapotranspiration comes from soil water?

Soil Moisture Conditions for Various Soil Textures



Depends on soil type


Available Water Capacity by Soil Texture

Textural Class	Available Water Capacity (Inches/Foot of Depth)	Available Water Capacity (mm/m)
Coarse sand	0.25–0.75	21-63
Fine sand	0.75–1.00	63-84
Loamy sand	1.10–1.20	92-100
Sandy loam	1.25–1.40	104-117
Fine sandy loam	1.50–2.00	125-167
Silt loam	2.00–2.50	167-208
Silty clay loam	1.80–2.00	150-167
Silty clay	1.50–1.70	125-141
Clay	1.20–1.50	100-125

What is my soil type?

What is my soil type?

Soil Textural Triangle



- Soil **texture** does not change
- Soil structure, porosity, water holding capacity, organic matter are affected by management:

Eg. 1% increase in SOM

- ≈ 5% increase water retention
- improved infiltration, structure

What is my soil type?

The screenshot displays the SoilWeb interface. The top navigation bar includes the UC Davis and NRCS logos. The main map area shows various soil units labeled with codes like YLA, RnA, CgD, W, DcE2, ReE, YrB, YmB, YnA, StF, JoF, AdA, HhF, and YrB. A red circle highlights a specific soil unit labeled 'YrB' on the map. The left sidebar provides detailed information for the selected unit:

Map Unit Name: **Yolo silt loam, 0 to 2 percent slopes** Symbol: **YsA**

Map Unit Composition

- 85% - **Yolo**
Geomorphic Position: *alluvial fans / Backslope*
- 15% - **Unnamed**
Horizon data *n/a*

Map Unit Data

- Map Unit Key: 459912
- Type: **Consociation** ?
- Farmland Class: *Prime farmland if irrigated*
- Available Water Storage (0-100cm): **17.6 cm**
- Max Flood Freq: *None*
- Drainage Class (Dominant Condition): **Well drained** ?
- Drainage Class (Wettest Component): **Well drained** ?
- Proportion of Hydric Soils: **0%** ?
- Min. Water Table Depth (Annual): *n/a*
- Min. Water Table Depth (April-June): *n/a*
- Min. Bedrock Depth: *n/a*
- Survey Metadata: *ca097 [NRCS Export: Sep 25 2014]*

At the bottom right, the coordinates are listed as Lat: 38.6278 and Lon: -122.8365. The Google logo is visible in the bottom left corner of the map area.

google: "soil web ucdavis"

What can I learn from knowing my soil type?

The screenshot displays the SoilWeb interface. The main map shows various soil units outlined in yellow, with a specific unit labeled 'YrX3' circled in red. The sidebar on the left provides detailed information for the selected unit, with several key attributes also circled in red.

Map Unit Name: *Yolo sandy loam, overwash, 0 to 5 percent slopes* Symbol: *YmB*

Map Unit Composition

- 85% - *Yolo*
Geomorphic Position: *alluvial fans / Backslope*
- 5% - *Cortina*
Horizon data *n/a* | [View Similar Data](#)
- 5% - *Zamora*
Horizon data *n/a* | [View Similar Data](#)
- 4% - *Pleasanton*
Horizon data *n/a* | [View Similar Data](#)
- 1% - *Unnamed*
Geomorphic Position: *flood plains*
Horizon data *n/a*

Map Unit Data

- Map Unit Key: 459908
- Type: *Consociation* ?
- Farmland Class: *Prime farmland if irrigated and either protected from flooding or not frequently flooded during the growing season*
- Available Water Storage (0-100cm): 15 cm
- Max Flood Freq: *Occasional*
- Drainage Class (Dominant Condition): *Well drained* ?
- Drainage Class (Wettest Component): *Well drained* ?
- Proportion of Hydric Soils: 1% ?
- Min. Water Table Depth (Annual): *n/a*

Map coordinates: Lat: 38.6266, Lon: -122.8204

Imagery ©2015, DigitalGlobe, USDA Farm Service Agency | 100 m | Terms of Use | Report a map error



Available Water Capacity by Soil Texture

Textural Class	Available Water Capacity (Inches/Foot of Depth)	Available Water Capacity (mm/m)
Coarse sand	0.25–0.75	21-63
Fine sand	0.75–1.00	63-84
Loamy sand	1.10–1.20	92-100
Sandy loam	1.25–1.40	104-117
Fine sandy loam	1.50–2.00	125-167
Silt loam	2.00–2.50	167-208
Silty clay loam	1.80–2.00	150-167
Silty clay	1.50–1.70	125-141
Clay	1.20–1.50	100-125

< Close

SoilWeb

Map Unit Name: **Yolo silt loam, 0 to 2 percent slopes** Symbol: **YsA**

Component Name: **Yolo**

Component Key: 11419299

[Official Series Description](#) | [Series Extent Explorer](#)

Soil Profiles

Typical Profile

Org. Matter ? >

Clay Sand

Ksat pH


Kr Factor EC

SAR CaCO₃

Gypsum

CEC @ pH7

Linear Ext.



[View Source Data](#)

Soil Taxonomy


Order: *Entisols*

Suborder: *Orthents* [Map of Suborders](#)

Greatgroup: *Xerorthents*

Subgroup: *Typic Xerorthents*

Family: *Fine-silt, mixed, nonacid, thermic Typic*



Field-based soil texture estimation methods

Classification in the field

A simple manual texture test is shown below.

Explanation:

- A ball about 2.5 cm diameter is formed from approximately 1 tablespoon of fine earth.
- Water is slowly dripped onto the soil until it approaches the sticky point, i.e., the point at which the soil just starts to stick to the hand.
- The extent to which the moist soil can be shaped by hand is indicative of its texture.

Textural class:

- (A) Sand - Soil remains loose and single-grained; can only be heaped into a pyramid.
- (B) Loamy sand - The soil contains sufficient silt and clay to become somewhat cohesive; can be shaped into a ball that easily falls apart.
- (C) Silt loam - Same as for loamy sand but can be shaped by rolling into a short, thick cylinder.
- (D) Loam - About equal sand, silt, and clay means the soil can be rolled into a cylinder about 15 cm long that breaks when bent.
- (E) Clay loam - As for loam, although soil can be bent into a U, but no further, without being broken.
- (F) Light clay - Soil can be bent into a circle that shows cracks.
- (G) Heavy clay - Soil can be bent into a circle without showing cracks.




Figure 4. Field method for evaluation of soil texture by feel. From Ilaco (1985).

NRCS:


Guide to Texture by Feel

Modified from S.J. Thien, 1979. A flow diagram for teaching texture by feel analysis. Journal of Agronomic Education, 8:54-55. (Click here for a high-resolution version of the graphic.)



http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/edu/?cid=nrcs142p2_054311

UCDavis:




<https://www.youtube.com/watch?v=GWZwbVJCNec>

Estimating depth of moisture, time to stress from soil type

Table 1. Percent moisture by volume

Texture	Perm. Wilting Point	Field Capacity	Available Water
	%		
Medium Sand	2.5	10.0	7.5
Fine Sand	3.3	12.5	9.2
Sandy Loam	5.0	16.7	11.7
Fine Sandy Loam	6.7	21.7	15.0
Loam	10.0	26.7	16.7
Silt Loam	11.7	29.2	17.5
Clay Loam	15.0	31.7	16.7
Clay	21.7	33.3	11.7



Meted.ucar.edu

For a loam soil $\approx 27\%$ moisture by volume at field capacity (FC).

- If the rooting depth is 20 inches, $20 \text{ in} \times 0.27 = 5.4$ inches of water by volume
- **Rule of thumb:** $FC/2 =$ available water. So, $5.4 / 2 = 2.7$ inches of available water.


When will stress begin?

- **Rule of thumb:** stress begins at available water / 2. So $2.7 / 2 = 1.35$ inches before stress.

Average daily ET in April in Windsor = 0.20 inches

- 1.35 inches / 0.20 inches per day
 - ≈ 7 days before stress.
 - ≈ 9 days if you use the table.

Soil type is a major factor in soil fertility



Nitrogen cycle ("Soil as a Plant Sees It", University of Nebraska, 1991).

Estimating soil fertility: Don't let the perfect be the enemy of the good



Figure 3. Nitrate-N quick hand test.




Figure 4. Nitrate color scale.

The Basic SOLVITA® Soil Response Color System

The patented gel-technology system indicates CO₂-respiration over a color range of 0 to 5 (see chart). In CO₂-Burst mode this corresponds to a range of 5 to 160 ppm CO₂-C. In BASAL mode it corresponds to a range of 0 – 55 ppm or 1 – 25 kg m²/year as CO₂.

All Solvita kits work with a basic visual color system, as shown below. By using the Solvita Digital Color Reader (DCR) the soil test values can be more accurately and precisely determined.



Sequence of Typical Soil Solvita Test Results:



http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_053274.pdf

Estimating soil fertility: Some rough rules of thumb

- Total N: Clay > Silt > Sand
 - effective/available N \neq total N
- ppm soil NO₃-N x 3.8 \approx lb N ac⁻¹ ft⁻¹ of soil
 - eg. 12 ppm NO₃-N (1st ft) x 3.8 \approx 46 lbs available N
- Manure: assume about 10% mineralized / season
 - 5 tons dry manure / acre; with 1.8% N \approx 18 lb ac⁻¹ yr⁻¹

Estimating soil fertility: Some rough rules of thumb

- Prior crop N contribution (depends on productivity of prior crop):
 - Tomato residue $\approx 25 \text{ lb ac}^{-1}$ returned
 - Alfalfa contribution $\approx 75 \text{ lb ac}^{-1} +$
- Soil organic matter (SOM) N mineralization:
 - $1.2\% \text{ OM } \% \times 30 \text{ lb N} / \% \text{ OM} \approx 36 \text{ lb ac}^{-1}$
- BUT! Mineralization rates vary (2-5%) year^{-1}
 - depend on C:N ratio, temperature, moisture, residue quality, etc.

Estimating soil fertility: Some rough rules of thumb

- Critical C:N ratio for residues ≈ 20
 - Residues with C:N > 20 will “fix” N

	C:N	Fix or release N?
Manure (Fresh)	15:1	RELEASE
Legumes (peas etc.)	15:1	RELEASE
Grass Clippings	20:1	EQUILIBRIUM
Weeds (Fresh)	25:1	FIX
Hay (Dry)	40:1	FIX
Weeds (Dry)	90:1	FIX
Straw, cornstalks	100:1	FIX
Sawdust	500:1	FIX

- Bottom line: fertility will be site-specific and a function of recent & long-term rotation/soil management

Estimating plant N removal from the system

If yield = **2500 lb acre⁻¹** and protein = **12.5%**

How much N am I removing from the system?

2500lb acre⁻¹ x 0.125 / 5.7 (protein to N factor)

- **≈ 55 lb N in grain**

55 x 1.33 (additional straw requirement)

- = 73 lb – 55lb
- **≈ 18 lb N in straw**

If we estimated that 100 lb of N are available from soil NO₃-N, SOM, and manure addition, will this crop experience N deficiency?




Figure 1-10 Leibig's Law of the Minimum states that the most limiting factor determines yield potential. Producers should minimize or eliminate the most limiting factor first, then the second most limiting factor, and so forth. Only in this manner can maximum yield potential be achieved (Source: Potash and Phosphate Institute).

Havlin, J. L. et al. 2005. Soil Fertility and Fertilizers, 7th ed.

Small grain growth stages and yield components




Image courtesy: University of Kentucky

plants acre⁻¹

tillers plant⁻¹

spikelets spike⁻¹

spike plant⁻¹

grains spikelet⁻¹

grains acre⁻¹

weight kernel⁻¹

GRAIN YIELD

Stand establishment

Seed selection

- Certified seed?
- plump uniform seeds, not cracked or broken, no weed seeds
 - < 1 year old
 - stored in dark, cool, dry conditions (free from pests and disease)



FAO. "Optimizing plant population, crop emergence and establishment"


Stand establishment

Germinability

- Rule of thumb: subtract 15% from germination test to get field germinability
- Seed-soil contact, soil moisture, temperature determine germination in the field
 - Rule of thumb: soil aggregates are no more than 3-4X seed size



Image courtesy: <http://www.bcg.org.au/>





FAO. "Optimizing plant population, crop emergence and establishment"


Stand establishment

Seeding Depth

- Rule of thumb: depth less than the coleoptile length.
 - coleoptile length varies
- Plant to moisture? or Moisture to follow?
 - Ensure sufficient moisture for complete germination



FAO. "Optimizing plant population, crop emergence and establishment"




FAO. "Optimizing plant population, crop emergence and establishment"

Stand establishment

Seeding rate

- Optimum plant populations and seeding rates require some empirical work. Range for most small grains: 75 – 125 lb acre⁻¹
 - spelt is higher (160 lb acre⁻¹ +)*
- Seeding rate interacts with weeds.
- BUT! For heirloom varieties, higher seeding rates will also tend to produce more lodging. WHY?
 - Site fertility also interacts with lodging outcomes.



FAO. "Optimizing plant population, crop emergence and establishment"

*<http://www.uvm.edu/extension/cropsoil/wp-content/uploads/TopTenGrowGrain.pdf>

Stand establishment

Seeding rate exercise:

- 2, 6 inch rows with 12 seeds per linear ft = 24 seeds / ft²
 - $43560 \text{ ft}^2 \text{ acre}^{-1} \times 24 \text{ seeds / ft}^2 = 1045440 \text{ seeds acre}^{-1}$
- For 10000 seeds lb⁻¹:
 - $1045440 \text{ seeds acre}^{-1} / 10000 \text{ seeds lb}^{-1}$
= 104 lb of seed acre⁻¹
- For 85% field germination rate:
 - $104 \text{ lb seed acre}^{-1} / 0.85 = 122 \text{ lb seed acre}^{-1}$

Stand establishment

Plant population exercise:

- 2, 6 inch rows with 12 seeds per linear ft = 24 seeds / ft²
 - $43560 \text{ ft}^2 \text{ acre}^{-1} \times 24 \text{ seeds / ft}^2 = 1045440 \text{ seeds acre}^{-1}$
- For 85% field germination rate:
 - $1045440 \text{ seed acre}^{-1} \times 0.85 = 888624 \text{ plants acre}^{-1}$
 - How many tillers per plant?
 - How many spikes per tiller?
 - How many spikelets per spike?
 - How many grains per spikelet?
 - Kernel weight?
- For 2500 lb acre⁻¹ yield, how many plants support 1 lb?
 - $888624 \text{ plants acre}^{-1} / 2500 \text{ lb acre}^{-1} = 355 \text{ plants lb}^{-1} \text{ grain}$
- When growing an heirloom variety, determining the optimum plant stand & seeding rate for your operation is worth the effort!

A brief note on TILLAGE

Why till?


- residue incorporation
- weed competition
- soil aeration
- aggregation (short term)
- seed-soil contact

Why no-till?

- reduced erosion, increased microbial diversity
- improved carbon retention, nutrient cycling (long term)
- improved water use efficiency, infiltration (long term)

Benefits of no-till are realized over the medium, long-term

- specialized equipment required
- short-term productivity losses
- residue, weed control are an issue for organic no-till



Variety Choice



Organic Seed Alliance

*Advancing the ethical development and stewardship
of the genetic resources of agricultural seed*

PO Box 772, Port Townsend, WA 98368

2013 California North Coast Organic Wheat Trials



In partnership with University of California
Cooperative Extension



With support from the California Wheat Commission
and Columbia Foundation

COLUMBIA FOUNDATION
Founded 1940

<http://seedalliance.org/publications>

Variety Choice

Variety	Type	Stripe rust	Relative maturity			Plant height			Lodging		Yield @ 13% moisture			Moisture		Test weight		Protein		
		CR	CR	FPF	ALL	CR	FPF	ALL	CR	FPF	CR	FPF	ALL	CR	FPF	CR	FPF	CR	FPF	ALL
		- % -	--- 1 to 9 scale ---			----- in -----			-- 1 to 9 scale --		----- lbs / acre -----			----- % -----		--- lbs / bu ---		----- % -----		
Yecora Rojo	HRS	76.7a	3.0e	2.8e	2.9e	18.0f	20.0f	19.0e	8.3bc	6.5abc	3265ab	1908a	2587a	13.1cd	11.7a	60.2c	59.6ab	12.6c	10.2e	11.4d
Canus	HRS	43.3b	5.0d	5.0d	5.0d	34.0c	30.0bc	32.0b	7.0c	3.0d	2609bc	1392b	2000b	13.1cd	10.8e	62.1b	59.9a	13.3b	13.5bc	13.4bc
Lassik	HRS	0.0c	4.8d	5.3d	5.0d	25.3d	24.0e	24.7d	8.8ab	8.0ab	3585a	1816a	2701a	12.5d	10.9de	63.9a	60.0a	12.7bc	12.6cd	12.7c
Red Fife	HRS	63.3a	7.3b	7.0b	7.1b	36.7b	30.5b	33.6b	8.0bc	6.3bc	1901de	616d	1258d	14.0ab	11.1cd	62b	59.9a	13.2b	14.2b	13.7b
Alturas	SWS	3.3c	5.0d	5.0d	5.0d	22.7e	26.3de	24.5d	7.0c	8.5a	2197cde	949c	1573cd	13.8bc	11.3bc	62.8b	58.1bc	11.1d	11.5de	11.3d
Diva	SWS	10.0c	6.0c	6.0c	6.0c	25.7d	27.5cd	26.6c	7.3c	5.75c	2465cd	896cd	1680c	13.9bc	11.2c	62.2b	60.2a	11.5d	11.5d	11.5d
Foisy	SWS	46.7b	9.0a	9.0a	9.0a	45.3a	35.3a	40.3a	9a	7.5abc	1692e	638cd	1165e	14.7a	11.5ab	60.8c	57.8c	15.2a	17.1a	16.2a
AVE		34.8	5.7	5.7	5.7	30	27.6	28.7	2.1	6.5	2531	1173	1852	13.6	11.2	62.0	59.4	12.8	12.9	12.9
CV (%)		25	5	5	5	6	7	6	45	22	23	18	23	3	2	1	2	3	7	5
LSD		12.7	0.4	0.4	0.3	2.5	3	1.9	1.4	2.1	791	314	411	0.76	0.3	0.7	1.7	0.5	1.3	0.7

CR = College of the Redwoods Farm

FPF = Front Porch Farm

ALL = Combined results from CR and FPF

Numbers in **bold** are the optimum greatest or least trait value or are not significantly different from the optimum

Letters after trait value indicate groups of varieties whose means are not significantly different for that trait.

Variety Choice

NORTHWEST CROPS & SOILS PROGRAM



2014 Heirloom Spring Wheat Seeding Rate Trial



Dr. Heather Darby, UVM Extension Agronomist
Erica Cummings, Katie Blair, Susan Monahan, Julian Post, Sara Ziegler
UVM Extension Crops and Soils Technicians
(802) 524-6501

Visit us on the web: <http://www.uvm.edu/extension/cropsoil>

© January 2015, University of Vermont Extension



The Whole Grain Connection

<http://www.sustainablegrains.org/>

List of available seeds

Common wheat (hexaploid, free threshing)
Triticum aestivum ssp aestivum

Variety (WGC catalog number) USDA accession number	Bearded or beardless	Historical notes (year collected by USDA or other)	Seed color (white or red)	Spring (short season) or winter (long season) type
Sonora (012) CItr 3036	beardless	Cultivar from landrace in Durango, Mexico. Perhaps the first successful wheat in Mexico from 1500. (1907)	Pale yellow (white)	Spring (shortest season)
Wit Wolkoring (013) PI 479660	beardless	Cultivar from South Africa. Presumed from landrace, but may be a cross. (1983)	Pale yellow (white)	Spring (shortest season)

<http://www.sustainablegrains.org/sitebuildercontent/sitebuilderfiles/wheatseedcatalog2015.pdf>

<http://www.uvm.edu/extension/cropsoil/grains>


Weed management

Tine weeding at early vegetative growth stage

- soil moisture, subsequent water will play into success/failure




First pass with the tine weeder after spring wheat emergence.



The work of the weeder

Weed management

Inter-seeding / relay-seeding red clover



Weed management


Stale seedbed techniques

- require precision in space and time



Image courtesy: <http://store.farmstart.ca/>

Weed management



Grants and Education to Advance Innovations in Sustainable Agriculture

[WebStore](#) | [Advanced Search](#) | [MySARE Login](#) | [SARE Social Media](#) | [SANET Listserv](#) | [Contact Us](#) | [Low Bandwidth](#) |

Search all SARE Sites

[Grants](#) | [Project Reports](#) | [Learning Center](#) | [Professional Development](#) | [State Programs](#) | [Events](#) | [Newsroom](#) | [About SARE](#)

Home » [Learning Center](#) » [Books](#) » [Steel in the Field](#) » [Text Version](#) » [Row Crop Tools](#) » [Cultivar, moderate-residue](#)

Font Size Print Share

You are reading the SARE book *Steel in the Field*.
[Order this publication.](#)

Text Version

[Acknowledgments](#)

[Publisher's Foreword](#)

[Cultivation In Context](#)

[How to Use This Book](#)

[Introduction to Tools for Agronomic Row Crops](#)

Row Crop Tools

- Rotary hoe, standard
- Rotary hoe, high-residue
- Rotary hoe accessories
- Flex-tine weeder
- Spike-tooth harrow

Cultivar, moderate-residue

Agronomic Row Crops Moderate-Residue Cultivator


Units intended for conservation tillage conditions (tilled residue with 30 percent coverage) or an untilled corn crop yielding up to 120 bushels/acre in loose to moderate soils with occasional stones up to 10 pounds.

Overview: These cultivators are a popular step up from a conventional cultivator for their ability to handle moderate weed pressure and moderate residue for less cost than a high-residue tool. S-tine tools have three to five shanks per row middle and generally weigh less than single-shank units. Choose sweep shape for your desired weed impact (i.e., bury, slice or uproot) and degree of soil/residue mixing. (See 'Cultivator sweep options,') Operating depth is about 1"

CROP height range estimate




WEED height range (annuals) estimate



Match tillage timing, depth and location to crop root growth. Weed control varies with soil conditions and weed density.

- Introduction To Cultivators
- Cultivator, low-residue
- **Cultivar, moderate-residue**
- Cultivator, high-residue
- Cultivator, maximum-residue
- Cultivator, rolling
- Cultivator, horizontal disk
- Cultivator sweeps, knives and wings
- Cultivator shields
- Cultivator components
- Hot-Tips for Flame Weeding
- Row-crop flamer
- Guidance Systems
- Guidance mirrors
- Guidance, furrower/wheel
- Guidance, ridge mechanical
- Hitch-steer guidance
- Side-shift guidance



Design Features: Compared with *low-residue* S-tine cultivators, these units usually have higher toolbar clearance (24" to 32") and longer front-to-back clearance (40" to 52"), allowing better flow between shanks, and between shanks and coulters, and overall stronger construction. Most have parallel linkage. Close-coupled, single-shank units are highly maneuverable and reduce

<http://www.sare.org/Learning-Center/Books/Steel-in-the-Field/Text-Version/Introduction-to-Tools-for-Agronomic-Row-Crops>

SARE Nationwide

Dual purpose wheat

Increased flexibility / risk mitigation for integrated crop-livestock systems

- generally fall-established wheat
- wait to graze until wheat is tillering
- grazing too long will severely reduce grain production
 - remove animals prior to first hollow stem




Figure 2. First hollow stem occurs when hollow stem equivalent to the diameter of a dime (1.5 cm) is present below the developing grain head.

Additional resources & other cool things



Southern Oregon Research and Extension Center

- Research
- Calendars
- Employee Resources
- Statewide Locations

Home » Small Farms

GROWING GRAINS ON A SMALL FARM

RESOURCES FOR PRODUCERS

Amber waves of grain are rippling across a handful of small farms in Southern Oregon. It could be a resurgence in regional production.

Eighty years ago, 41 varieties of wheat were grown on a million acres in Oregon. Small farms grew wheat. Today, there is much less wheat, and what is grown is mostly produced on the large commodity scale, as on the bigger farms in the Klamath basin.

Since small farms in Southwestern Oregon no longer produce much locally consumed grain, almost a generation of knowledge and infrastructure has been lost. The local foods movement, and the niche marketing opportunities it affords, have led to producers' renewed interest in growing grains for the local market.

In January, 2010, a group of bakers, brewers, chefs, millers and farmers met at OSU Extension to collaborate on ways to meet consumer demand for locally-produced grain products. This meeting was the first in a series of seven classes entitled, "Growing Grains on a Small Farm." The series, funded by a USDA Sustainable Agriculture Research and Education (SARE) grant, covered all aspects of grain production throughout the growing season. Each class took place on a farm in Southern Oregon and included a tour of the grain operation, as well as presentations by producers and university specialists.

The resources and links on this page are the results of that class series, as well as data from informal wheat trialing performed during the 2010 growing season.



BCS Tiller with Sickle Bar Mower at the Bjorn Everson Farm

General Information on Growing Grains	Livestock Feed	Economics
Direct Marketing	Wheat Trial Data	Equipment
Resources for Homesteader		

- [SOREC Home](#)
- [Faculty & Staff](#)
- [Contact Us](#)
- [About Our Center](#)
- [Find Us](#)
- [Employment Opportunities at SOREC](#)

What We Offer

- [4-H Youth](#)
- [4-H FourThoughts Newsletter Jan. - March 2015](#)
- [Natural Resources Youth Program](#)
- [Master Gardeners-Home Gardening](#)
- [Family and Community Health](#)
- [Viticulture & Enology](#)
- [Small Farms](#)
 - [OSU Small Farms News](#)
 - [Small Farms Class Proceedings](#)
 - [Small Farms Resources](#)
 - [Growing Grains-Resources](#)
- [Livestock & Forage](#)
- [Land Steward Program](#)
- [Commercial Agriculture](#)
- [Forestry and Natural Resources](#)
- [Citizen Fire Academy Program](#)
- [Recycling](#)

Social Media

- [College of Agricultural Science social media](#)
- [Jackson County 4-H](#)
- [Oregon State University Small Farms](#)
- [Southern Oregon Research & Extension Center](#)

Additional resources & other cool things

ational

x

www.tillersinternational.org/index.html



"Encouraging an Attitude of Experimentation to Produce More Local Food with Less Global Fuel"

Home About Donations Gallery Classes Events Store News Contact

- Oxen & Draft Horses
- Farming
- International Rural Development
- Blacksmithing & Metal Work
- Woodworking & Construction
- Food & Fiber
- Museum
- TechGuides
- e-News Sign-up
- Smallmouth CD



"To preserve, study, and exchange low-capital technologies that increase the sustainability and productivity of people in rural communities"


<http://www.tillersinternational.org/>

Additional resources & other cool things

Small Grain Harvesting Cl... x

← → ↻ <https://www.youtube.com/watch?v=hxNIeLUgYpg>

YouTube MUSIC AWARDS 🔍



0:24 / 0:47

Small Grain Harvesting Class at Tillers International

<https://www.youtube.com/watch?v=hxNIeLUgYpg>

Additional resources & other cool things

it no-till seed dr x

farmhack.org/tools/home-built-no-till-seed-drill

DONATE

FARM HACK

A COMMUNITY FOR FARM INNOVATION

TOOLS CONVERSATIONS CALENDAR HOST AN EVENT SHOPS PEOPLE

Login Register

Home built no-till seed drill

Short description:

This is my home built no-till seed drill I use for planting legume mix winter cover crop seed in our stone fruit orchard. It produces very uniform seed spacing and depth with minimal soil disturbance, allowing optimum germination in our dry farmed environment.



<http://farmhack.org/tools/home-built-no-till-seed-drill>

Additional resources & other cool things

powered thresher, x

farmhack.org/tools/bicycle-powered-thresherfan-mill-and-dehuller-1

FARM HACK  A COMMUNITY FOR FARM INNOVATION



TOOLS CONVERSATIONS CALENDAR HOST AN EVENT SHOPS PEOPLE Login Register

bicycle powered thresher, fan mill, and dehuller

Short description:

prototypes of a suite of bike powered tools for small scale grain processing; a thresher, a fan mill/winnower, and a dehuller. Many thanks to Olaf B-N for making this video:

<https://www.youtube.com/watch?v=Lgnmhtbgyfg>



<http://farmhack.org/tools/bicycle-powered-thresherfan-mill-and-dehuller-1>

Hope springs eternal...



Thank you!

contact: melundy@ucanr.edu

