HEALTHY ENVIRONMEN

HEALTHY COMMUNITIES • HEALTHY CALIFORNIAN

University of California

Agriculture and Natural Resources

Making a Difference for California



UCCE/DWR Weekly Crop Water Use Report

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C)

04/12/19 through 04/18/19

Crops (Leafout Date)	#148 Merced			 #39 Parlier			#258 Lemon Cove			
	4/12 - 4/18	Accum'd	4/19 - 4/25	4/12 - 4/18	Accum'd	4/19 - 4/25	4/12 - 4/18	Accum'd	4/19 - 4/25	
	Water	Seasonal	Estimated	Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc	 Use	Water Use	ETc	Use	Water Use	ETc	1
Almonds (3/10) *	0.92	3.57	0.96	0.98	3.94	0.96	0.96	3.68	0.94	
Pistachio (4/19) * **	0.00	0.00	0.07	0.00	0.00	0.07	0.00	0.00	0.07	
Citrus (2/1)	0.88	5.76	0.89	0.93	6.25	0.91	0.92	5.71	0.87	
Raisin Grapes (3/16) (11 ft. row spacing) ***	0.22	0.54	0.26	0.23	0.57	0.28	0.23	0.54	0.24	
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis) ***	0.29	0.81	0.33	0.31	0.93	0.35	0.30	0.88	0.31	
Walnuts (4/15)	0.08	0.08	0.14	0.09	0.09	0.14	0.08	0.08	0.14	
Stone Fruit (3/20)	0.37	1.08	0.47	0.39	1.22	0.47	0.38	1.10	0.47	
Past 7 days precipitation (inches)		0.02			0.12			0.19		
Accumulated precipitation (inches) (1/1/2019)		8.82			5.78			8.68		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

*** Raisin Grapes and Winegrapes Irrigation should hold off until midday leaf water potential drops to -1.0 MPa, before that soil moisture reservoir is sufficient to supply the vine water demand. Update will be sent shortly once the county wide leaf water potential reaches approximate -1.0 MPa. Growers should adjust the irrigation start date based on the individual vineyard location and soil type.

PA	ST WEEK	LY APPLII	ED WATE	R IN INCHE	ES, ADJUST	FED FOR E	FFICIENC	Y 1					
Crops	#148 Merced					#39 Parlier	•		#258 Lemon Cove				
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/10)	1.4	1.2	1.1	1.0	1.5	1.3	1.2	1.0	1.5	1.3	1.1	1.0	
Pistachio (4/19)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Citrus (2/1)	1.4	1.2	1.0	0.9	1.4	1.2	1.1	1.0	1.4	1.2	1.1	1.0	
Raisin Grapes (3/16) (11 ft. row spacing)***	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis) ***	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Walnuts (4/15)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
Stone Fruit (3/20)	0.6	0.5	0.4	0.4	0.6	0.5	0.5	0.4	0.6	0.5	0.4	0.4	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST V	VEEKLY A	PPLIED W	ATER IN G	GALLON P	ER TREE C	R VINE					
Crops		#148 Merc		#39 Parlier			#258 Lemon Cove					
Almonds 115 Trees/A	331	283	260	236	354	307	283	236	354	307	260	236
Pistachio 106 Trees/A	0	0	0	0	0	0	0	0	0	0	0	0
Citrus 110 Trees/A	346	296	247	222	346	296	272	247	346	296	272	247
Raisin Grapes 566 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Winegrapes 622 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Walnuts 76 Trees/A	36	36	36	36	36	36	36	36	36	36	36	36
Stonefruit 172 Trees/A	95	79	63	63	95	79	79	63	95	79	63	63
For further information concerning all counties receiving this report, contact	the Fresno	Co. Farm Ad	visor's office	at (559) 24	1-7526.							

University of **California** Agriculture and Natural Resources



UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

												(
	WEEKLY SOIL MOISTURE LOSS IN INCHES											
(Estimated Crop Evapotranspiration or ET _C)												
04/12/19 through 04/18/19												
Crops (Leafout Date) #124 Panoche #2 Five Points #15 Stratford												
Crops (Learout Date)	4/12 - 4/18	Accum'd	4/19 - 4/25		4/12 - 4/18	Accum'd	4/19 - 4/25		4/12 - 4/18	Accum'd	4/19 - 4/25	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated		Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc		Use	Water Use	ETc	
Almonds (3/10) *	1.04	3.98	1.13		1.13	4.67	1.17		1.06	4.28	1.17	
Pistachio (4/19) * **	0.00	0.00	0.10		0.00	0.00	0.12		0.00	0.00	0.12	
Citrus (2/1)	0.99	6.29	1.03		1.08	7.41	1.07		1.00	6.55	1.07	
Raisin Grapes (3/16) (11 ft. row spacing) ***	0.25	0.60	0.31		0.28	0.71	0.33		0.27	0.66	0.33	
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis) ***	0.34	0.98	0.40		0.38	1.17	0.42		0.34	1.03	0.42	
Walnuts (4/15)	0.09	0.09	0.19		0.11	0.11	0.21		0.09	0.09	0.21	
Stone Fruit (3/20)	0.41	1.22	0.57		0.44	1.41	0.59		0.42	1.35	0.59	
Past 7 days precipitation (inches)		0.05				0.23				0.35		
Accumulated precipitation (inches) (1/1/2019)		4.03				5.45				4.14		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

*** Raisin Grapes and Winegrapes Irrigation should hold off until midday leaf water potential drops to -1.0 MPa, before that soil moisture reservoir is sufficient to supply the vine water demand. Update will be sent shortly once the county wide leaf water potential reaches approximate -1.0 MPa. Growers should adjust the irrigation start date based on the individual vineyard location and soil type.

PA	ST WEEK	LY APPLI	ED WATEF	R IN INCHE	ES, ADJUST	FED FOR E	FFICIENC	Y ¹				
Crops		#124 Pano	che			#2 Five Poi	ints					
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/10)	1.6	1.4	1.2	1.1	1.7	1.5	1.3	1.2	1.6	1.4	1.2	1.1
Pistachio (4/19)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Citrus (2/1)	1.5	1.3	1.2	1.0	1.7	1.4	1.3	1.1	1.5	1.3	1.2	1.1
Raisin Grapes (3/16) (11 ft. row spacing) ***	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis) ***	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Walnuts (4/15)	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Stone Fruit (3/20)	0.6	0.5	0.5	0.4	0.7	0.6	0.5	0.5	0.6	0.6	0.5	0.4

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST V	VEEKLY A	PPLIED W	ATER IN (GALLON P	ER TREE (DR VINE					
Crops		#124 Pano	#15 Stratfo	15 Stratford								
Almonds 115 Trees/A	378	331	283	260	401	354	307	283	378	331	283	260
Pistachio 106 Trees/A	0	0	0	0	0	0	0	0	0	0	0	0
Citrus 110 Trees/A	370	321	296	247	420	346	321	272	370	321	296	272
Raisin Grapes 566 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Winegrapes 622 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Walnuts 76 Trees/A	36	36	36	36	71	36	36	36	36	36	36	36
Stonefruit 172 Trees/A	95	79	79	63	111	95	79	79	95	95	79	63
For further information concerning all counties receiving this report, contact	t the Fresno	Co. Farm Ad	visor's office	at (559) 24	11-7526.				-			