University of **California** Agriculture and Natural Resources



UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 04/19/19 through 04/25/19

Crops (Leafout Date)	#148 Merced			#39 Parlier			#258 Lemon Cove			
	4/19 - 4/25	Accum'd	4/26 - 5/2	4/19 - 4/25	Accum'd	4/26 - 5/2	4/19 - 4/25	Accum'd	4/26 - 5/2	
	Water	Seasonal	Estimated	Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc	Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (3/10) *	1.06	4.63	1.05	1.14	5.08	1.11	1.11	4.79	1.04	
Pistachio (4/19) * **	0.10	0.10	0.14	0.12	0.12	0.14	0.10	0.20	0.14	
Citrus (2/1)	0.98	6.74	0.93	1.07	7.32	0.99	1.04	6.75	0.92	
Raisin Grapes (3/16) (11 ft. row spacing) ***	0.38	0.92	0.42	0.40	0.97	0.42	0.38	0.92	0.42	
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis) ***	0.44	1.25	0.44	0.45	1.38	0.49	0.44	1.33	0.43	
Walnuts (4/15)	0.17	0.25	0.56	0.20	0.29	0.59	0.18	0.26	0.55	
Stone Fruit (3/20)	0.54	1.62	0.51	0.58	1.80	0.56	0.55	1.65	0.50	
Past 7 days precipitation (inches)		0.00			0.00			0.00		
Accumulated precipitation (inches) (1/1/2019)		8.82			5.78			8.68		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY 1

Crops		#148 Merc	ed			#39 Parlier			#258 Lemon Cove				
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/10)	1.6	1.4	1.2	1.1	1.8	1.5	1.3	1.2	1.7	1.5	1.3	1.2	
Pistachio (4/19)	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	
Citrus (2/1)	1.5	1.3	1.2	1.0	1.6	1.4	1.3	1.1	1.6	1.4	1.2	1.1	
Raisin Grapes (3/16) (11 ft. row spacing)***	0.6	0.5	0.4	0.4	0.6	0.5	0.5	0.4	0.6	0.5	0.4	0.4	
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis) ***	0.7	0.6	0.5	0.5	0.7	0.6	0.5	0.5	0.7	0.6	0.5	0.5	
Walnuts (4/15)	0.3	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.2	
Stone Fruit (3/20)	0.8	0.7	0.6	0.6	0.9	0.8	0.7	0.6	0.8	0.7	0.6	0.6	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

	IASIV	EEKLI A	I LLED W	AIEKIN	ALLONI	EK TKEE (IX VIIIE						
Crops		#148 Merce	ed			#39 Parlier			#258 Lemon Cove				
Almonds 115 Trees/A	378	331	283	260	425	354	307	283	401	354	307	283	
Pistachio 106 Trees/A	50	25	25	25	50	25	25	25	50	25	25	25	
Citrus 110 Trees/A	370	321	296	247	395	346	321	272	395	346	296	272	
Raisin Grapes 566 Vines/A	29	24	19	19	29	24	24	19	29	24	19	19	
Winegrapes 622 Vines/A	31	26	22	22	31	26	22	22	31	26	22	22	
Walnuts 76 Trees/A	107	71	71	71	107	107	71	71	107	71	71	71	
Stonefruit 172 Trees/A	126	111	95	95	142	126	111	95	126	111	95	95	

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

^{***} Raisin Grapes and Winegrapes Irrigation should hold off until midday leaf water potential drops to -1.0 MPa, before that soil moisture reservoir is sufficient to supply the vine water demand. Update will be sent shortly once the county wide leaf water potential reaches approximate -1.0 MPa. Growers should adjust the irrigation start date based on the individual vineyard location and soil type.

HEALTHY FOOD SYSTEMS • HEALTHY ENVIRONMENTS • HEALTHY COMMUNITIES • HEALTHY CALIFORNIANS

University of California Agriculture and Natural Resources



UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 04/19/19 through 04/25/19

Crops (Leafout Date)	#124 Panoche			#2 Five Points				#15 Stratford			
	4/19 - 4/25	Accum'd	4/26 - 5/2	4/19 - 4/25	Accum'd	4/26 - 5/2		4/19 - 4/25	Accum'd	4/26 - 5/2	
	Water	Seasonal	Estimated	Water	Seasonal	Estimated		Water	Seasonal	Estimated	İ
	Use	Water Use	ETc	Use	Water Use	ETc		Use	Water Use	ETc	
Almonds (3/10) *	1.24	5.22	1.29	1.30	5.97	1.31		1.24	5.52	1.29	
Pistachio (4/19) * **	0.12	0.12	0.14	0.14	0.14	0.15		0.12	0.12	0.14	
Citrus (2/1)	1.16	7.45	1.13	1.21	8.62	1.19		1.16	7.71	1.16	
Raisin Grapes (3/16) (11 ft. row spacing) ***	0.46	1.07	0.51	0.47	1.18	0.55		0.45	1.11	0.53	
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis) ***	0.52	1.50	0.55	0.54	1.71	0.56		0.51	1.54	0.56	
Walnuts (4/15)	0.20	0.29	0.68	0.21	0.32	0.71		0.20	0.29	0.70	
Stone Fruit (3/20)	0.62	1.84	0.63	0.68	2.09	0.64		0.62	1.97	0.63	
Past 7 days precipitation (inches)		0.00			0.07		-		0.00		
Accumulated precipitation (inches) (1/1/2019)		4.03			5.52				4.14		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

^{***} Raisin Grapes and Winegrapes Irrigation should hold off until midday leaf water potential drops to -1.0 MPa, before that soil moisture reservoir is sufficient to supply the vine water demand. Update will be sent shortly once the county wide leaf water potential reaches approximate -1.0 MPa. Growers should adjust the irrigation start date based on the individual vineyard location and soil type.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹													
Crops	#124 Panoche					#2 Five Poi	nts		#15 Stratford				
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/10)	1.9	1.7	1.5	1.3	2.0	1.7	1.5	1.4	1.9	1.7	1.5	1.3	
Pistachio (4/19)	0.2	0.2	0.1	0.1	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.1	
Citrus (2/1)	1.8	1.5	1.4	1.2	1.9	1.6	1.4	1.3	1.8	1.5	1.4	1.2	
Raisin Grapes (3/16) (11 ft. row spacing) ***	0.7	0.6	0.5	0.5	0.7	0.6	0.6	0.5	0.7	0.6	0.5	0.5	
Winegrapes (3/16) (10 ft. spacing on California Sprawl Trellis) ***	0.8	0.7	0.6	0.5	0.8	0.7	0.6	0.6	0.8	0.7	0.6	0.5	
Walnuts (4/15)	0.3	0.3	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.3	0.2	0.2	
Stone Fruit (3/20)	1.0	0.8	0.7	0.7	1.0	0.9	0.8	0.7	1.0	0.8	0.7	0.7	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE												
Crops		#124 Panoo	che			#2 Five Poi	nts					
Almonds 115 Trees/A	449	401	354	307	472	401	354	331	449	401	354	307
Pistachio 106 Trees/A	50	50	25	25	50	50	50	25	50	50	25	25
Citrus 110 Trees/A	444	370	346	296	469	395	346	321	444	370	346	296
Raisin Grapes 566 Vines/A	34	29	24	24	34	29	29	24	34	29	24	24
Winegrapes 622 Vines/A	35	31	26	22	35	31	26	26	35	31	26	22
Walnuts 76 Trees/A	107	107	71	71	107	107	71	71	107	107	71	71
Stonefruit 172 Trees/A	158	126	111	111	158	142	126	111	158	126	111	111
For further information concerning all counties receiving this report, contact	the Fresno C	Co. Farm Ad	visor's office	at (559) 24	1-7526.							

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.