

---

# Fungicide Control of Phomopsis Cane and Leaf Spot on Grapevine: 2015 Field Trial

---

W. Douglas Gubler, Trang T. Nguyen and Nicholas S. Morris

Department of Plant Pathology, University of California, Davis, CA 95616.

---

Department of Plant Pathology,  
University of California Cooperative Extension,  
University of California, Davis – October 2015

---

Published 2015 at: [http://plantpathology.ucdavis.edu/Cooperative\\_Extension/](http://plantpathology.ucdavis.edu/Cooperative_Extension/)  
Copyright © 2015 by the Regents of the University of California, Davis campus. All Rights Reserved.

## Introduction

Phomopsis cane and leaf spot of grapevine is caused by the fungal pathogen, *Phomopsis viticola*. On leaves, the disease is manifest as tiny dark spots with yellow halos on the leaf blade and veins (Bettiga *et al.*, 2014). Similar spotting can occur on petioles or on the basal portion of infected shoots, and heavy infection on shoots may cause a scab-like appearance (Bettiga *et al.*, 2014). Spring rains occurring after budbreak stimulate spore release, dispersal and infection (Gubler *et al.*, 2008). Spores released from overwintering pycnidia on canes and spurs are spread by rain to young shoots. Infection occurs most readily when moisture remains on the green tissue for many hours (Nita *et al.*, 2008).

In California, the disease can be economically important during wet years in the north coast and in the northern San Joaquin Valley. Economic loss is generally minor, except during years when damage to shoots and fruit reduces the number of fruit clusters. In the northern San Joaquin Valley, susceptible grape varieties include Cabernet Sauvignon, Pinot Noir, Chardonnay, Thompson Seedless, and Grenache (Flaherty *et al.*, 1992). Fungicide applications are made during the spring months to protect shoots.

A field trial was conducted at the UC Davis Plant Pathology Farm in northeastern Solano County, CA to evaluate the efficacy of registered fungicides on control of Phomopsis cane and leaf spot. The variety Thompson Seedless was used in this study.

## Materials and Methods

The trial was conducted on 3 rows of vines in a 34 year-old Thompson Seedless vineyard (8ft vine x 12 ft row spacing), using a complete randomized design with 4 replicates. Plots consisted of 4 adjacent vines. Fungicides were applied with a handgun sprayer (Nifty-Fifty circulating tank) using 75 gallons/acre on 3 Mar and 100 gallons/acre on 17 Mar. At the time of the first application, shoots were roughly 20 cm in length.

Disease was assessed on 19 Aug by rating disease severity on each vine by counting the number of infected shoots. Trial models were analyzed using the ANOVA Tests for data; P-values for trial was  $P < 0.05$ . Means comparisons were made using Student's t-test with  $\alpha = 0.05$ .


**Table 1.** Experimental Design

Experimental design	Complete randomized design with 4 replicates.		
Experimental unit	4 vines = 1 plot		
Plot area	384 ft <sup>2</sup> (row spacing = 12 ft, vine spacing = 8 ft)		
Area/treatment	1536 ft <sup>2</sup> (4 reps = 1 treatment)	Area/treatment	0.035 acre/treatment
Volume water/acre	75 gallons = 2.63 gal/4 replicates (3/3/2015) 100 gallons = 3.5 gal/4 replicates (3/17/2015)		
Application method	Handgun sprayers (attached to Nifty Fifty brand 25 or 50 gallon sprayers).		

**Table 2.** Treatments examined in the trial. FP = formulated product.

Flag	Product(s)	FP/Acre	FP/4 replicate plots
W	Unsprayed control	none	none
BKS	Merivon + Dyneamic	6.5 fl oz + 0.25%	6.7 ml + 26.5 ml at 75 gal or 34.1 ml at 100 gal
YKC	Captan Gold	2.5 lbs	39.7 g
OKD	Sovran + Dyneamic	4 oz + 0.25%	4.0 g + 26.5 ml at 75 gal or 34.1 ml at 100 gal


Figure 1. Map.




## Results

Several rain events from the beginning of March to early May (Figure 2) provided wetness for Phomopsis infection on emerging green tissue. Overall disease pressure was low, however all treatments significantly decreased disease compared to untreated control.

Figure 2. Daily precipitation data for Davis, California from 1 Mar to 19 Aug 2015. Data from CIMIS station 6 (<http://www.cimis.water.ca.gov/>).



**Figure 3.** Daily temperature data for Davis, California from 1 Mar to 19 Aug 2015. Data from CIMIS station 6 (<http://www.cimis.water.ca.gov/>).



**Table 3.** Treatment effects on shoot disease symptoms. Treatment means with different letters are significantly different according to Student's t-test ( $\alpha = 0.05$ ). '\*' Treatment followed by rate per acre.

Treatment	Mean # of infected shoots
Sovran, 4 oz + Dyneamic, 0.25% (v/v)*	1.13 c
Captan Gold, 2.5 lbs	1.19 bc
Merivon, 6.5 fl oz + Dyneamic, 0.25% (v/v)	1.94 b
Unsprayed control	3.69 a

## References

Grape Pest Management. UC DANR Publication 3343, 3<sup>rd</sup> edition. Regents of the University of California. Bettiga *et al.*, 2014

Gubler WD, Smith RJ, Varela LJ, Vasquez S, Stapleton JJ, and Purcell AH. (2008) UC IPM Management Guidelines: Grape, UCANR Publication 3448, Diseases.

Nita M, Ellis MA, and Madden LV (2008) Variations in disease incidence of Phomopsis cane and leaf spot of grape in commercial vineyards in Ohio. *Plant Disease* 92:1053-1061.

## Appendix: Materials

Product	Active ingredient(s) and concentration	Class	Manufacturer
Captan Gold 80WDG	captan (78.2%)	Pthalamide (M4)	Adama
Dyne-Amic	polyalkyleneoxide modified polydimethylsiloxane, nonionic emulsifiers, methyl ester of C16-C18 fatty acids (99%)	Adjuvant	Helena Chemical Co.
Merivon	fluxopyroxad (21%), pyraclostrobin (21%)	SDHI (7)/QoI (11)	BASF
Sovran	kresoxim-methyl (50%)	QoI (11)	Cheminova

*Appendix references:* (1) Adaskaveg, et al. 2012. Efficacy and timing of fungicides, bactericides and biologicals for deciduous tree fruit, nut, strawberry, and vine crops 2012, available at <http://www.ipm.ucdavis.edu/PDF/PMG/fungicideefficacytiming.pdf>, (2) 2014 Fungicide trials, available at [http://plantpathology.ucdavis.edu/Cooperative\\_Extension/](http://plantpathology.ucdavis.edu/Cooperative_Extension/), (3) various sources including product labels and/or MSDS.