
Final Report: Evaluation of Biotam (*Trichoderma asperellum* & *Trichoderma gamsii*) & crab life powder as Pruning Wound Protectants Against Selected Fungi Associated with Grapevine Trunk Diseases

R. Blundell, M. Lynch, T. Haden, M. Arreguin, T. Gallagher and A. Eskalen
Department of Plant Pathology, University of California, Davis, CA, 95616

University of California Cooperative Extension,
Department of Plant Pathology,
University of California, Davis, January 2020

Report Summary

Grapevine trunk diseases (GTDs) represent a major threat to the future economic sustainability of table grapes and wine grapes. Several taxonomically unrelated groups of Ascomycete fungi cause trunk diseases in grapevines including *Eutypa lata* and *Neofusicoccum parvum*. (1). Following precipitation events, fungal spores (sexual and asexual) become airborne and colonize exposed wood vessels caused by pruning. Total disease control is virtually unattainable because of the huge number of wounds made on an individual grapevine and extended period of wound susceptibility but one mitigation practice is to apply a protectant to exposed pruning wounds (2, 3, 4, 5).

This trial was conducted at the UC Davis Plant Pathology Fieldhouse Facility (38.522591, -121.760719) from March to September 2019. Treatments were a randomized block design in an 8 year old Sauvignon Blanc vineyard.

Materials and Methods

A. Experimental design

Experimental design	Complete randomized block design
Experimental unit	2 spurs of each = 1 plot x 10 rep
Plot area	110 ft ² (row spacing = 11 ft, vine spacing = 7 ft)
Application method	Hand held spray bottle

B. Experimental treatments

The treatments described in this report were conducted for experimental purposes only and crops treated in a similar manner may not be suitable for commercial or other use.

Flag	Treatment Name	Application rate	FP/10 vines (0.6L spray bottle)	Date applied
B	Crab life powder	0.5 lb/100 ga	0.35 g	After pruning
GS	Biotam	2 lb/A	1.43 g	After pruning
BC	Biotam & crab life powder	2 lb/A & 0.5 lb/100 ga	1.43 g + 0.35 g	After pruning
OKD	Rally + Topsin M + organosilicone surfactant	2.25 oz + 1.25 lbs/A	0.05 g + 0.45 g + 12 µl	After pruning
W	Untreated control	N/A	N/A	N/A

C. Map

[illegible]

D. Application calendar

		March
Flag	Treatment Name	21
B	Crab life powder	x
GS	Biotam	x
BC	Biotam + Crab life powder	x
OKD	Rally + Topsin M + organosilicone surfactant	x
W	Untreated control	N/A

E. Vine Management

During the application period, vines were irrigated by drip irrigation. Sucker shoot removal and leafing were done during the duration of trial.

F. Data Collection and Statistics

The efficacy of the treatments controlling the GTDs were recorded as the Mean Percentage of Infection (MPI). This was calculated by: (Number of GTD infected samples/Number of total samples) x 100. There were a total of 20 repetitions per GTD per treatment. Treatments were compared against the untreated control and a standard control. Means comparisons were made using Dunnett's test $\alpha=0.05$.

Results




Figure 1. Evaluation of pruning wound treatments mean percent infection (MPI) rates with *E. lata* located at UC Davis Plant Pathology Field House, 2019. Bars represent the least mean square of percent infection. Bars with a different letter are different according to Dunnett's test ($p = 0.05$).




Figure 2. Evaluation of pruning wound treatments mean percent infection (MPI) rates with *N. parvum* located at UC Davis Plant Pathology Field House, 2019. Bars represent the least mean square of percent infection. Bars with a different letter are different according to Dunnett's test ($p = 0.05$).

Acknowledgements

Thanks to the various industry donors for providing testing materials. Thanks to Bryan Pellissier and Lexi Sommers-Miller for their field support.

Appendix

Flag color	Trade Name	Active Ingredient	Manufacturer
B	Crab life powder		
GS	Biotam	Trichoderma asperellum & Trichoderma gamsii	ISAGRO-USA
BC	Biotam & Crab life powder	Trichoderma asperellum & Trichoderma gamsii	ISAGRO-USA

Literature Cited

1. Moller, W.J., and A.N. Kasimatis. 1978. Dieback of grapevines caused by *Eutypa armeniacae*. Plant Dis. Rep. 62:254-258.
2. Eskalen, A., A.J. Feliciano, and W.D. Gubler. 2007. Susceptibility of grapevine pruning wounds and symptom development in response to infection by *Phaeoacremonium aleophilum* and *Phaeomoniella chlamydospora*. Plant Dis. 91:1100-1104.
3. Petzoldt, C.H., M.A. Sall, and W.J. Moller. 1983. Factors determining the relative number of ascospores released by *Eutypa armeniacae* in California. Plant Dis. 67:857-860.
4. Rooney-Latham, S., A. Eskalen, and W.D. Gubler. 2005. Occurrence of *Togninia minima* perithecia in esca-affected vineyards in California. Plant Dis. 89:867-871.
5. Urbez-Torres, J.R., and W.D. Gubler. 2008. Double pruning, a potential method to control Bot canker disease of grapes, and susceptibility of grapevine pruning wounds to infection by Botryosphaeriaceae. Abstr. Phytopathol. Mediterr. 48:185.