University of California Agriculture and Natural Resources



UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 05/20/22 through 05/26/22

Crops (Leafout Date)	#148 Merced				#39 Parlier		#258 Lemon Cove			
	5/20- 5/26	Accum'd	5/27 - 6/2	5/20- 5/26	Accum'd	5/27 - 6/2	5/20- 5/26	Accum'd	5/27 - 6/2	
	Water	Seasonal	Estimated	Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc	Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (3/1) *	1.78	12.43	1.65	1.78	12.68	1.67	1.74	12.33	1.62	
Pistachio (4/8) * **	1.75	6.25	1.77	1.75	6.42	1.79	1.71	6.29	1.73	
Citrus (2/1)	1.32	13.30	1.16	1.31	13.53	1.18	1.29	13.21	1.13	
Raisin Grapes (3/15) (11 ft. row spacing)	1.16	4.56	1.15	1.18	4.68	1.16	1.15	4.56	1.13	
Winegrapes (3/15) (10 ft. spacing on California Sprawl Trellis) ***	1.10	4.68	1.10	1.08	4.77	1.11	1.08	4.70	1.07	
Walnuts (4/8)	1.62	7.99	1.52	1.62	8.28	1.54	1.58	8.12	1.49	
Stone Fruit (3/10)	1.38	7.32	1.36	1.38	7.42	1.38	1.35	7.29	1.33	
Past 7 days precipitation (inches)		0.00			0.00			0.00		
Accumulated precipitation (inches) (1/1/2022)		1.88			1.79			1.82		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED	FOR EFFICIENCY 1
---	------------------

Crops	#148 Merced					#39 Parlier			#258 Lemon Cove			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	2.7	2.4	2.1	1.9	2.7	2.4	2.1	1.9	2.7	2.3	2.0	1.8
Pistachio (4/8)	2.7	2.3	2.1	1.8	2.7	2.3	2.1	1.8	2.6	2.3	2.0	1.8
Citrus (2/1)	2.0	1.8	1.6	1.4	2.0	1.7	1.5	1.4	2.0	1.7	1.5	1.4
Raisin Grapes (3/15) (11 ft. row spacing)***	As	sume all gr	ape	1.2	Assume all grape 1			1.2	Assume all grape			1.2
Winegrapes (3/15) (10 ft. spacing on California Sprawl Trellis) ***	irriga	ation type is	s drip	1.2	irrigation type is drip			1.1	irrigation type is drip			1.1
Walnuts (4/8)	2.5	2.2	1.9	1.7	2.5	2.2	1.9	1.7	2.4	2.1	1.9	1.7
Stone Fruit (3/10)	2.1	1.8	1.6	1.5	2.1	1.8	1.6	1.5	2.1	1.8	1.6	1.4

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops	#148 Merced					#39 Parlier			#258 Lemon Cove			
Almonds 115 Trees/A	638	567	496	449	638	567	496	449	638	543	472	425
Pistachio 106 Trees/A	673	573	523	448	673	573	523	448	648	573	498	448
Citrus 110 Trees/A	494	444	395	346	494	420	370	346	494	420	370	346
Raisin Grapes 566 Vines/A	Assume all grape			58	Assume all grape 58			Assume all grape			58	
Winegrapes 622 Vines/A	irriga	ation type is	s drip	52	irrigation type is drip 48			irrigation type is drip			48	
Walnuts 76 Trees/A	893	786	679	607	893	786	679	607	857	750	679	607
Stonefruit 172 Trees/A	332	284	253	237	332	284	253	237	332	284	253	221
For further information concerning all counties receiving this report, contact	the Fresno C	Co. Farm Ad	visor's office	at (559) 24	1-7526.							

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

University of California Agriculture and Natural Resources



UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 05/20/22 through 05/26/22

Crops (Leafout Date)	#124 Panoche				#2 Five Points			#15 Stratford			
	5/20 - 5/26	Accum'd	5/27 - 6/2		5/20 - 5/26	Accum'd	5/27 - 6/2	5/20 - 5/26	Accum'd	5/27 - 6/2	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (3/1) *	2.11	14.26	1.82		2.08	14.90	1.90	2.12	15.10	1.90	
Pistachio (4/8) * **	2.05	7.08	1.95		2.02	7.33	2.03	2.06	7.45	2.03	
Citrus (2/1)	1.52	14.88	1.26		1.50	15.66	1.32	1.53	15.55	1.32	
Raisin Grapes (3/15) (11 ft. row spacing)	1.34	5.08	1.25		1.33	5.30	1.31	1.36	5.37	1.31	
Winegrapes (3/15) (10 ft. spacing on California Sprawl Trellis)	1.27	5.24	1.19		1.24	5.43	1.24	1.27	5.54	1.24	
Walnuts (4/8)	1.90	9.07	1.65		1.88	9.45	1.73	1.91	9.62	1.73	
Stone Fruit (3/10)	1.62	8.22	1.47		1.59	8.56	1.55	1.62	8.65	1.55	
Past 7 days precipitation (inches)		0.00		_		0.00		 _	0.00		
Accumulated precipitation (inches) (1/1/2022)		1.11				1.11			1.30		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY 1												
Crops	#124 Panoche					#2 Five Poi	ints					
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	3.2	2.8	2.5	2.2	3.2	2.8	2.4	2.2	3.3	2.8	2.5	2.2
Pistachio (4/8)	3.2	2.7	2.4	2.2	3.1	2.7	2.4	2.1	3.2	2.7	2.4	2.2
Citrus (2/1)	2.3	2.0	1.8	1.6	2.3	2.0	1.8	1.6	2.4	2.0	1.8	1.6
Raisin Grapes (3/15) (11 ft. row spacing)	As	sume all gr	ape	1.4	Assume all grape			1.4	Assume all grape			1.4
Winegrapes (3/15) (10 ft. spacing on California Sprawl Trellis)	irriga	ation type is	s drip	1.3	irriga	ation type is	s drip	1.3	irrigation type is drip			1.3
Walnuts (4/8)	2.9	2.5	2.2	2.0	2.9	2.5	2.2	2.0	2.9	2.5	2.2	2.0
Stone Fruit (3/10)	2.5	2.2	1.9	1.7	2.4	2.1	1.9	1.7	2.5	2.2	1.9	1.7

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE													
Crops		#124 Pano	che			#2 Five Poi	ints						
Almonds 115 Trees/A	756	661	590	519	756	661	567	519	779	661	590	519	
Pistachio 106 Trees/A	797	673	598	548	772	673	598	523	797	673	598	548	
Citrus 110 Trees/A	568	494	444	395	568	494	444	395	592	494	444	395	
Raisin Grapes 566 Vines/A	As	sume all gr	ape	67	Assume all grape 67			67	Assume all grape			67	
Winegrapes 622 Vines/A	irriga	ation type is	s drip	57	irrigation type is drip			57	irrigation type is drip		s drip	57	
Walnuts 76 Trees/A	1036	893	786	715	1036	893	786	715	1036	893	786	715	
Stonefruit 172 Trees/A	395	347	300	268	379	332	300	268	395	347	300	268	
For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.													

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.