☐ UCCE/DWR Weekly Crop Water Use Report





WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 07/22/22 through 07/28/22

Crops (Leafout Date)	7	#148 Merced			#39 Parlier			#258 Lemon Cove			
	7/22- 7/28	Accum'd	7/29- 8/4		7/22- 7/28	Accum'd	7/29- 8/4	7/22- 7/28	Accum'd	7/29- 8/4	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (3/1) *	2.10	30.65	1.96		2.12	31.25	1.89	1.93	29.38	1.89	
Pistachio (4/8) * **	2.15	25.48	2.03		2.19	26.06	1.96	1.98	24.37	1.96	
Citrus (2/1)	1.28	24.81	1.19		1.29	25.32	1.12	1.16	24.00	1.12	
Raisin Grapes (3/15) (11 ft. row spacing)	1.40	16.96	1.31		1.43	17.35	1.25	1.28	16.18	1.25	
Winegrapes (3/15) (10 ft. spacing on California Sprawl Trellis)	1.60	18.32	1.49		1.62	18.70	1.43	1.46	17.44	1.43	
Walnuts (4/8)	2.22	26.18	2.10		2.26	26.84	2.03	2.05	25.16	2.03	
Stone Fruit (3/10)	2.14	23.75	2.03		2.18	24.19	1.96	1.96	22.58	1.96	
Past 7 days precipitation (inches)		0.00				0.00			0.00		
Accumulated precipitation (inches) (1/1/2022)		1.98				1.84			1.82		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENC	\mathbf{Y}^{1}
---	------------------

That weeker hiteleb which is inches, housafed for efficiency												
Crops		#148 Merc	ed			#39 Parlier	•		#258 Lemon Cove			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	3.2	2.8	2.5	2.2	3.3	2.8	2.5	2.2	3.0	2.6	2.3	2.0
Pistachio (4/8)	3.3	2.9	2.5	2.3	3.4	2.9	2.6	2.3	3.0	2.6	2.3	2.1
Citrus (2/1)	2.0	1.7	1.5	1.3	2.0	1.7	1.5	1.4	1.8	1.5	1.4	1.2
Raisin Grapes (3/15) (11 ft. row spacing)	As	Assume all grape			Assume all grape 1.5			1.5	Assume all grape			1.4
Winegrapes (3/15) (10 ft. spacing on California Sprawl Trellis)	irrig	irrigation type is drip		1.7	irrigation type is drip		1.7	irrigation type is drip		s drip	1.5	
Walnuts (4/8)	3.4	3.0	2.6	2.3	3.5	3.0	2.7	2.4	3.2	2.7	2.4	2.2
Stone Fruit (3/10)	3.3	2.9	2.5	2.3	3.4	2.9	2.6	2.3	3.0	2.6	2.3	2.1

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops			#39 Parlier	•		#258 Lemon Cove						
Almonds 115 Trees/A	756	661	590	519	779	661	590	519	708	614	543	472
Pistachio 106 Trees/A	822	722	623	573	847	722	648	573	747	648	573	523
Citrus 110 Trees/A	494	420	370	321	494	420	370	346	444	370	346	296
Raisin Grapes 566 Vines/A	Assume all grape			72	Ass	Assume all grape 72			Assume all grape			67
Winegrapes 622 Vines/A	irrigation type is drip			74	irrigation type is drip 74			74	irrigation type is drip			65
Walnuts 76 Trees/A	1215	1072	929	822	1251	1072	965	857	1143	965	857	786
Stonefruit 172 Trees/A	521	458	395	363	537	458	410	363	474	410	363	332
For further information concerning all counties receiving this report, contact	the Fresno (Co. Farm Ad	visor's offic	e at (559) 24	41-7526.							

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

HEALTHY FOOD SYSTEMS • HEALTHY ENVIRONMENTS • HEALTHY COMMUNITIES • HEALTHY CALIFORNIANS

University of California Agriculture and Natural Resources Making a Difference for California



UCCE/DWR Weekly Crop Water Use Report

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 07/22/22 through 07/28/22

Crops (Leafout Date)	#	124 Panoch	e	#2 Five Points				#15 Stratford			
	7/22 - 7/28	Accum'd	7/29 - 8/4	7/22 - 7/28	Accum'd	7/29 - 8/4		7/22 - 7/28	Accum'd	7/29 - 8/4	
	Water	Seasonal	Estimated	Water	Seasonal	Estimated		Water	Seasonal	Estimated	
	Use	Water Use	ETc	Use	Water Use	ETc		Use	Water Use	ETc	
Almonds (3/1) *	2.26	34.41	1.96	2.29	35.53	2.10		2.36	36.20	2.10	
Pistachio (4/8) * **	2.33	28.20	2.03	2.35	28.97	2.17		2.41	29.53	2.17	
Citrus (2/1)	1.40	27.58	1.19	1.40	28.58	1.26		1.44	28.79	1.26	
Raisin Grapes (3/15) (11 ft. row spacing)	1.50	18.71	1.31	1.52	19.30	1.43		1.57	19.67	1.43	
Winegrapes (3/15) (10 ft. spacing on California Sprawl Trellis)	1.73	20.15	1.49	1.76	20.68	1.61		1.81	21.11	1.61	
Walnuts (4/8)	2.40	29.06	2.10	2.43	29.96	2.24		2.49	30.57	2.24	
Stone Fruit (3/10)	2.31	26.32	2.03	2.34	27.07	2.17		2.40	27.58	2.17	
Past 7 days precipitation (inches)		0.00			0.00	<u>-</u>	_		0.00	_	
Accumulated precipitation (inches) (1/1/2022)		1.23			1.11				1.30		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

PA	PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY 1												
Crops	#124 Panoche					#2 Five Po	ints		#15 Stratford				
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/1)	3.5	3.0	2.7	2.4	3.5	3.1	2.7	2.4	3.6	3.1	2.8	2.5	
Pistachio (4/8)	3.6	3.1	2.7	2.5	3.6	3.1	2.8	2.5	3.7	3.2	2.8	2.5	
Citrus (2/1)	2.2	1.9	1.6	1.5	2.2	1.9	1.6	1.5	2.2	1.9	1.7	1.5	
Raisin Grapes (3/15) (11 ft. row spacing)	As	Assume all grape			Assume all grape 1.6			1.6	Assume all grape			1.7	
Winegrapes (3/15) (10 ft. spacing on California Sprawl Trellis)	irrigation type is drip		1.8	irriga	irrigation type is drip		1.9	irrigation type is drip		s drip	1.9		
Walnuts (4/8)	3.7	3.2	2.8	2.5	3.7	3.2	2.9	2.6	3.8	3.3	2.9	2.6	
Stone Fruit (3/10)	3.6	3.1	2.7	2.4	3.6	3.1	2.8	2.5	3.7	3.2	2.8	2.5	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST WEEKLY	APPLIED	WATER	IN GALLON	PER TREE OR VINE
--	-------------	---------	-------	-----------	------------------

Crops			#2 Five Poi	ints		#15 Stratford						
Almonds 115 Trees/A	826	708	638	567	826	732	638	567	850	732	661	590
Pistachio 106 Trees/A	897	772	673	623	897	772	698	623	922	797	698	623
Citrus 110 Trees/A	543	469	395	370	543	469	395	370	543	469	420	370
Raisin Grapes 566 Vines/A	Assume all grape			77	Ass	Assume all grape 77			As	sume all gra	ape	82
Winegrapes 622 Vines/A	irrigation type is drip			79	irrigation type is drip 83			83	irrigation type is drip			83
Walnuts 76 Trees/A	1322	1143	1000	893	1322	1143	1036	929	1358	1179	1036	929
Stonefruit 172 Trees/A	568	489	426	379	568	489	442	395	584	505	442	395
For further information concerning all counties receiving this report, contact	the Fresno (Co. Farm Ad	visor's office	e at (559) 2	41-7526.							

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.